

*Application  
for  
United States Letters Patent*

To all whom it may concern:

Be it known that **Arlindo L. Castelhana, et al.,**

have invented certain new and useful improvements in

**Compounds Specific To Adenosine A<sub>3</sub> Receptor And Uses Thereof**

of which the following is a full, clear and exact description.

**COMPOUNDS SPECIFIC TO ADENOSINE A<sub>3</sub> RECEPTOR AND USES THEREOF**

This application claims the benefit of U.S. Provisional Application No. 60/169,036, filed December 2, 1999.

5

Throughout this application, various publications are referenced by author and date. Full citations for these publications may be found throughout the specification. The disclosures of these publications in their entirety are hereby incorporated by reference into this application in order to more fully describe the state of the art as known to those skilled therein as of the date of the invention described and claimed herein.

15 **Background of the Invention**

Adenosine is an ubiquitous modulator of numerous physiological activities, particularly within the cardiovascular and nervous systems. The effects of adenosine appear to be mediated by specific cell surface receptor proteins. Adenosine modulates diverse physiological functions including induction of sedation, vasodilation, suppression of cardiac rate and contractility, inhibition of platelet aggregability, stimulation of gluconeogenesis and inhibition of lipolysis. In addition to its effects on adenylyate cyclase, adenosine has been shown to open potassium channels, reduce flux through calcium channels, and inhibit or stimulate phosphoinositide turnover through receptor-mediated mechanisms (See for example, C.E. Muller and B. Stein "Adenosine Receptor Antagonists: Structures and Potential Therapeutic Applications," *Current Pharmaceutical Design*, 2:501 (1996) and C.E. Muller "A<sub>1</sub>-Adenosine Receptor Antagonists," *Exp. Opin. Ther. Patents* 7(5):419 (1997)).

Adenosine receptors belong to the superfamily of purine receptors which are currently subdivided into P<sub>1</sub> (adenosine) and P<sub>2</sub> (ATP, ADP, and other nucleotides) receptors. Four receptor subtypes for the nucleoside adenosine have been cloned so far from various species including humans. Two receptor subtypes (A<sub>1</sub> and A<sub>2a</sub>) exhibit affinity for adenosine

in the nanomolar range while two other known subtypes  $A_{2b}$  and  $A_3$  are low-affinity receptors, with affinity for adenosine in the low-micromolar range.  $A_1$  and  $A_3$  adenosine receptor activation can lead to an inhibition of adenylate cyclase activity, while  $A_{2a}$  and  $A_{2b}$  activation causes a stimulation of adenylate cyclase.

A few  $A_1$  antagonists have been developed for the treatment of cognitive disease, renal failure, and cardiac arrhythmias. It has been suggested that  $A_{2a}$  antagonists may be beneficial for patients suffering from Morbus Parkinson (Parkinson's disease). Particularly in view of the potential for local delivery, adenosine receptor antagonists may be valuable for treatment of allergic inflammation and asthma. Available information (for example, Nyce & Metzger "DNA antisense Therapy for Asthma in an Animal Model" *Nature* (1997) 385: 721-5) indicates that in this pathophysiologic context,  $A_1$  antagonists may block contraction of smooth muscle underlying respiratory epithelia, while  $A_{2b}$  or  $A_3$  receptor antagonists may block mast cell degranulation, mitigating the release of histamine and other inflammatory mediators.  $A_{2b}$  receptors have been discovered throughout the gastrointestinal tract, especially in the colon and the intestinal epithelia. It has been suggested that  $A_{2b}$  receptors mediate cAMP response (Strohmeier et al., *J. Bio. Chem.* (1995) 270:2387-94).

Adenosine receptors have also been shown to exist on the retinas of various mammalian species including bovine, porcine, monkey, rat, guinea pig, mouse, rabbit and human (See, Blazynski et al., *Discrete Distributions of Adenosine Receptors in Mammalian Retina*, *Journal of Neurochemistry*, volume 54, pages 648-655 (1990); Woods et al., *Characterization of Adenosine  $A_1$ -Receptor Binding Sites in Bovine Retinal Membranes*, *Experimental Eye Research*, volume 53, pages 325-331 (1991); and Braas et al., *Endogenous adenosine and adenosine receptors localized to ganglion cells*

5 Nucleoside Transport Sites in a Cultured Human Retinal Cell  
Line Established By SV-40 T Antigen Gene, Current Eye  
Research, volume 13, pages 109-118 (1994)).

Compounds which regulate the uptake of adenosine uptake have previously been suggested as potential therapeutic agents for the treatment of retinal and optic nerve head damage. In U.S. Patent No. 5,780,450 to Shade, Shade discusses the use of adenosine uptake inhibitors for treating eye disorders. Shade does not disclose the use of specific A<sub>2</sub> receptor inhibitors. The entire contents of U.S. Patent No. 5,780,450 are hereby incorporated herein by reference.

Additional adenosine receptor antagonists are needed as pharmacological tools and are of considerable interest as drugs for the above-referenced disease states and/or conditions.

### Summary of the Invention


The present invention is based on compounds which selectively bind to adenosine A<sub>3</sub> receptor, thereby treating a disease associated with A<sub>3</sub> adenosine receptor in a subject by administering to the subject a therapeutically effective amount of such compounds. The disease to be treated are associated with, for example, asthma, hypersensitivity, rhinitis, hay fever, serum sickness, allergic vasculitis, atopic dermatitis, dermatitis, psoriasis, eczema, idiopathic pulmonary fibrosis, eosinophilic cholecystitis, chronic airway inflammation, hypereosinophilic syndromes, eosinophilic gastroenteritis, edema, urticaria, eosinophilic myocardial disease, episodic angioedema with eosinophilia, inflammatory bowel disease, ulcerative colitis, allergic granulomatosis, carcinomatosis, eosinophilic granuloma, familial histiocytosis, hypertension, mast cell degranulation, tumor, cardiac hypoxia, cerebral ischemia, diuresis, renal failure, neurological disorder, mental disorder, cognitive disorder, myocardial ischemia, bronchoconstriction, arthritis, autoimmune disease, Crohn's disease, Grave's disease, diabetes, multiple sclerosis, anaemia, psoriasis, fertility disorders, lupus erythematosus, reperfusion injury, brain arteriole diameter, the release of allergic mediators, scleroderma, stroke, global ischemia, central nervous system disorder, cardiovascular disorder, renal disorder, inflammatory disorder, gastrointestinal disorder, eye disorder, allergic disorder, respiratory disorder, or immunological disorder.

The present invention is based, at least in part, on the discovery that certain N-6 substituted 7-deazapurines, described *infra*, can be used to treat a N-6 substituted 7-deazapurine responsive state. Examples of such states include those in which the activity of the adenosine receptors is increased, e.g., bronchitis, gastrointestinal disorders, or asthma. These states can be characterized in that adenosine receptor activation can lead to the inhibition

or stimulation of adenylate cyclase activity. Compositions and methods of the invention include enantiomerically or diastereomerically pure N-6 substituted 7-deazapurines. Preferred N-6 substituted 7-deazapurines include those which  
5 have an acetamide, carboxamide, substituted cyclohexyl, e.g., cyclohexanol, or a urea moiety attached to the N-6 nitrogen through an alkylene chain.

The present invention pertains to methods for modulating an  
10 adenosine receptor(s) in a mammal by administering to the mammal a therapeutically effective amount of a N-6 substituted 7-deazapurine, such that modulation of the adenosine receptor's activity occurs. Suitable adenosine  
15 receptors include the families of A<sub>1</sub>, A<sub>2</sub>, or A<sub>3</sub>. In a preferred embodiment, the N-6 substituted 7-deazapurine is a adenosine receptor antagonist.

The invention further pertains to methods for treating N-6  
substituted 7-deazapurine disorders, e.g., asthma,  
20 bronchitis, allergic rhinitis, chronic obstructive pulmonary disease, renal disorders, gastrointestinal disorders, and eye disorders, in a mammal by administering to the mammal a therapeutically effective amount of a N-6 substituted 7-  
25 deazapurine, such that treatment of the disorder in the mammal occurs. Suitable N-6 substituted 7 deazapurines include those illustrated by the general formula I:




(I)

and pharmaceutically acceptable salts thereof.  $R_1$  and  $R_2$  are each independently a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety or together form a substituted or unsubstituted heterocyclic ring.  $R_3$  is  
5 a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety.  $R_4$  is a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety.  $R_5$  and  $R_6$  are each independently a halogen atom, e.g., chlorine, fluorine, or bromine, a hydrogen atom or a substituted or unsubstituted  
10 alkyl, aryl, or alkylaryl moiety or  $R_4$  and  $R_5$  or  $R_5$  and  $R_6$  together form a substituted or unsubstituted heterocyclic or carbocyclic ring.

In certain embodiments,  $R_1$  and  $R_2$  can each independently be  
15 a substituted or unsubstituted cycloalkyl or heteroarylalkyl moieties. In other embodiments,  $R_3$  is a hydrogen atom or a substituted or unsubstituted heteroaryl moiety. In still other embodiments,  $R_4$ ,  $R_5$  and  $R_6$  can each be independently a heteroaryl moieties. In a preferred embodiment,  $R_1$  is a  
20 hydrogen atom,  $R_2$  is a cyclohexanol, e.g., trans-cyclohexanol,  $R_3$  is phenyl,  $R_4$  is a hydrogen atom,  $R_5$  is a methyl group and  $R_6$  is a methyl group. In still another embodiment,  $R_1$  is a hydrogen atom,  $R_2$  is

25



$R_3$  is phenyl,  $R_4$  is a hydrogen atom and  $R_5$  and  $R_6$  are methyl  
30 groups.

The invention further pertains to pharmaceutical compositions for treating a N-6 substituted 7-deazapurine responsive state in a mammal, e.g., asthma, bronchitis, allergic rhinitis,  
35 chronic obstructive pulmonary disease, renal disorders, gastrointestinal disorders, and eye disorders. The pharmaceutical composition includes a therapeutically

effective amount of a N-6 substituted 7-deazapurine and a pharmaceutically acceptable carrier.

007027 9493260  
The present invention also pertains to packaged  
5 pharmaceutical compositions for treating a N-6 substituted 7-deazapurine responsive state in a mammal. The packaged pharmaceutical composition includes a container holding a therapeutically effective amount of at least one N-6  
10 substituted 7-deazapurine and instructions for using the N-6 substituted 7-deazapurine for treating a N-6 substituted 7-deazapurine responsive state in a mammal.

The invention further pertains to compounds of formula I wherein

15  $R_1$  is hydrogen;  
 $R_2$  is substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkyl, or  $R_1$  and  $R_2$  together form a substituted or unsubstituted heterocyclic ring;  
 $R_3$  is unsubstituted or substituted aryl;  
20  $R_4$  is hydrogen; and  
 $R_5$  and  $R_6$  are each independently hydrogen or alkyl, and pharmaceutically acceptable salts thereof. The deazapurines of this embodiment may advantageously be selective  $A_3$  receptor antagonists. These compounds may be useful for numerous  
25 therapeutic uses such as, for example, the treatment of asthma, kidney failure associated with heart failure, and glaucoma. In a particularly preferred embodiment, the deazapurine is a water soluble prodrug that is capable of being metabolized in vivo to an active drug by, for example,  
30 esterase catalyzed hydrolysis.

In yet another embodiment, the invention features a method for inhibiting the activity of an adenosine receptor (e.g.,  $A_3$ ) in a cell, by contacting the cell with N-6 substituted 7-deazapurine (e.g., preferably, an adenosine receptor  
35 antagonist).



In another aspect, the invention features a method for treating damage to the eye of an animal (e.g., a human) by administering to the animal an effective amount of an N-6 substituted 7-deazapurine of formula I. Preferably, the N-6 substituted 7-deazapurine is an antagonist of A<sub>3</sub> adenosine receptors in cells of the animal. The damage is to the retina or the optic nerve head and may be acute or chronic. The damage may be the result of, for example, glaucoma, edema, ischemia, hypoxia or trauma.


10

The invention also features a pharmaceutical composition comprising a N-6 substituted compound of formula I. Preferably, the pharmaceutical preparation is an ophthalmic formulation (e.g., an periocular, retrobulbar or intraocular injection formulation, a systemic formulation, or a surgical irrigating solution).

15

In yet another embodiment, the invention features a compound having the formula II:

20



25

30

(II)

wherein X is N or CR<sub>6</sub>; R<sub>1</sub> and R<sub>2</sub> are each independently hydrogen, or substituted or unsubstituted alkoxy, aminoalkyl, alkyl, aryl, or alkylaryl, or together form a substituted or unsubstituted heterocyclic ring, provided that both R<sub>1</sub> and R<sub>2</sub> are both not hydrogen; R<sub>3</sub> is substituted or unsubstituted alkyl, arylalkyl, or aryl; R<sub>4</sub> is hydrogen or substituted or unsubstituted C<sub>1</sub>- C<sub>6</sub> alkyl; L is hydrogen, substituted or unsubstituted alkyl, or R<sub>4</sub> and L together form a substituted or unsubstituted heterocyclic or carbocyclic ring; R<sub>6</sub> is hydrogen, substituted or unsubstituted alkyl, or halogen; Q is CH<sub>2</sub>, O, S, or NR<sub>7</sub>, wherein R<sub>7</sub> is hydrogen or substituted or unsubstituted C<sub>1</sub>- C<sub>6</sub> alkyl; and W is unsubstituted or substituted alkyl, cycloalkyl, aryl, arylalkyl, biaryl, heteroaryl, substituted carbonyl, substituted thiocarbonyl, or substituted sulfonyl; provided that if R<sub>3</sub> is pyrrolidino, then R<sub>4</sub> is not methyl. The invention also pertains to pharmaceutically acceptable salts and prodrugs of the compounds of the invention.


In an advantageous embodiment, X is CR<sub>6</sub> and Q is CH<sub>2</sub>, O, S, or NH in formula II, wherein R<sub>6</sub> is as defined above.

In another embodiment of formula II, X is N.  
The invention further pertains to a method for inhibiting the activity of an adenosine receptor (e.g., an A<sub>2b</sub> adenosine receptor) in a cell by contacting the cell with a compound of the invention. Preferably, the compound is an antagonist of the receptor.

The invention also pertains to a method for treating a gastrointestinal disorder (e.g., diarrhea) or a respiratory disorder (e.g., allergic rhinitis, chronic obstructive pulmonary disease) in an animal by administering to an animal an effective amount of a compound of formula II (e.g., an

antagonist of A<sub>2c</sub>). Preferably, the animal is a human.

This invention also features a compound having the structure:



wherein R<sub>1</sub> is H and R<sub>2</sub> is cyclopropyl methylamino carbonylethyl, *cis*-3-hydroxy cyclopentyl, acetamido butyl, methylamino carbonylamino butyl, ethylamino carbonylamino propyl, methylamino carbonylamino propyl, 2-acetyl amino-3-methyl butyl, N,N-diethylamino carbonylamino ethyl, thioacetamido ethyl, 3-amino acetyloxy cyclopentyl, 3-hydroxy cyclopentyl, 2-pyrrolyl carbonyl aminoethyl, 2-imidazolidinone ethyl, 1-aminocarbonyl-2-methyl propyl, 1-aminocarbonyl-2-phenyl ethyl, 3-hydroxy azetidino, 2-imidazolyl ethyl, acetamido ethyl, 1-(R)-phenyl-2-hydroxyethyl, N-methylaminocarbonyl pyridyl-2-methyl, or R<sub>1</sub>, R<sub>2</sub> and the nitrogen together are 3-acetamido piperadino, 3-hydroxy pyrrolidino, 3-methyloxy carbonylmethyl pyrrolidino, 3-aminocarbonylmethyl pyrrolidino, or 3-hydroxymethyl piperadino.

wherein R<sub>3</sub> is a substituted or unsubstituted four to six membered ring, pyrrole, thiophene, furan, thiazole, imidazole, pyrazole, 1,2,4-triazole, pyridine, 2(1H)-pyridone, 4(1H)-pyridone, pyrazine, pyrimidine, pyridazine, isothiazole, isoxazole, oxazole, tetrazole,

naphthalene, tetralin, naphthyridine, benzofuran,  
benzothiophene, indole, 2,3-dihydroindole, 1H-indole,  
indoline, benzopyrazole, 1,3-benzodioxole, benzoxazole,  
purine, coumarin, chromone, quinoline,  
5 tetrahydroquinoline, isoquinoline, benzimidazole,  
quinazoline, pyrido[2,3-b]pyrazine, pyrido[3,4-  
b]pyrazine, pyrido[3,2-c]pyridazine, purido[3,4-b]-  
pyridine, 1H-pyrazole[3,4-d]pyrimidine, pteridine,  
2(1H)-quinolone, 1(2H)-isoquinolone, 1,4-benzisoxazine,  
10 benzothiazole, quinoxaline, quinoline-N-oxide,  
isoquinoline-N-oxide, quinoxaline-N-oxide, quinazoline-  
N-oxide, benzoxazine, phthalazine, or cinnoline.

wherein R<sub>5</sub> is H, alkyl, substituted alkyl, or cycloalkyl;  
15 wherein R<sub>6</sub> is H, alkyl, substituted alkyl, aryl, or  
substituted aryl.

This invention also features a method for inhibiting the  
20 activity of an A<sub>3</sub> adenosine receptor in a cell, which  
comprises contacting said cell with the above-mentioned  
compounds.

25

Detailed Description

The features and other details of the invention will now be more particularly described and pointed out in the claims. It will be understood that the particular embodiments of the invention are shown by way of illustration and not as limitations of the invention. The principle features of this invention can be employed in various embodiments without departing from the scope of the invention.

- 10 The present invention pertains to methods for treating a N-6 substituted 7-deazapurine responsive state in a mammal. The methods include administration of a therapeutically effective amount of a N-6 substituted 7-deazapurine, described *infra*, to the mammal, such that treatment of the N-6 substituted 7-  
15 deazapurine responsive state in the mammal occurs.

The language "N-6 substituted 7-deazapurine responsive state" is intended to include a disease state or condition characterized by its responsiveness to treatment with a N-6 substituted 7-deazapurine of the invention as described *infra*, e.g., the treatment includes a significant  
20 diminishment of at least one symptom or effect of the state achieved with a N-6 substituted 7-deazapurine of the invention. Typically such states are associated with an increase of adenosine within a host such that the host often  
25 experiences physiological symptoms which include, but are not limited to, release of toxins, inflammation, coma, water retention, weight gain or weight loss, pancreatitis, emphysema, rheumatoid arthritis, osteoarthritis, multiple organ failure, infant and adult respiratory distress  
30 syndrome, allergic rhinitis, chronic obstructive pulmonary disease, eye disorders, gastrointestinal disorders, skin tumor promotion, immunodeficiency and asthma. (See for example, C.E. Muller and B. Stein "Adenosine Receptor  
35 Antagonists: Structures and Potential Therapeutic Applications," *Current Pharmaceutical Design*, 2:501 (1996) and C.E. Muller "A<sub>1</sub>-Adenosine Receptor Antagonists," *Exp.*

007007:91932460

Opin. Ther. Patents 7(5):419 (1997) and I. Feoktistove, R. Polosa, S. T. Holgate and I. Biaggioni "Adenosine A<sub>2B</sub> receptors: a novel therapeutic target in asthma?" *TIPS* 19; 148 (1998)). The effects often associated with such symptoms include, but are not limited to, fever, shortness of breath, nausea, diarrhea, weakness, headache, and even death. In one embodiment, a N-6 substituted 7-deazapurine responsive state includes those disease states which are mediated by stimulation of adenosine receptors, e.g., A<sub>1</sub>, A<sub>2a</sub>, A<sub>2b</sub>, A<sub>3</sub>, etc., such that calcium concentrations in cells and/or activation of PLC (phospholipase C) is modulated. In a preferred embodiment, a N-6 substituted 7-deazapurine responsive state is associated with adenosine receptor(s), e.g., the N-6 substituted 7-deazapurine acts as an antagonist. Examples of suitable responsive states which can be treated by the compounds of the invention, e.g., adenosine receptor subtypes which mediate biological effects, include central nervous system (CNS) effects, cardiovascular effects, renal effects, respiratory effects, immunological effects, gastro-intestinal effects and metabolic effects. The relative amount of adenosine in a subject can be associated with the effects listed below; that is increased levels of adenosine can trigger an effect, e.g., an undesired physiological response, e.g., an asthmatic attack.

CNS effects include decreased transmitter release (A<sub>1</sub>), sedation (A<sub>1</sub>), decreased locomotor activity (A<sub>2a</sub>), anticonvulsant activity, chemoreceptor stimulation (A<sub>2</sub>) and hyperalgesia. Therapeutic applications of the inventive compounds include treatment of dementia, Alzheimer's disease and memory enhancement.

Cardiovascular effects include vasodilation (A<sub>2a</sub>), (A<sub>2b</sub>) and (A<sub>3</sub>), vasoconstriction (A<sub>1</sub>), bradycardia (A<sub>1</sub>), platelet inhibition (A<sub>2a</sub>), negative cardiac inotropy and dromotropy (A<sub>1</sub>), arrhythmia, tachycardia and angiogenesis. Therapeutic

applications of the inventive compounds include, for example, prevention of ischaemia-induced impairment of the heart and cardiotonics, myocardial tissue protection and restoration of cardiac function.

5

Renal effects include decreased GFR ( $A_1$ ), mesangial cell contraction ( $A_1$ ), antidiuresis ( $A_1$ ) and inhibition of renin release ( $A_1$ ). Suitable therapeutic applications of the inventive compounds include use of the inventive compounds as  
10 diuretic, natriuretic, potassium-sparing, kidney-protective/prevention of acute renal failure, antihypertensive, anti-oedematous and anti-nephritic agents.

Respiratory effects include bronchodilation ( $A_2$ ),  
15 bronchoconstriction ( $A_1$ ), chronic obstructive pulmonary disease, allergic rhinitis, mucus secretion and respiratory depression ( $A_2$ ). Suitable therapeutic applications for the compounds of the invention include anti-asthmatic applications, treatment of lung disease after transplantation  
20 and respiratory disorders.

Immunological effects include immunosuppression ( $A_2$ ), neutrophil chemotaxis ( $A_1$ ), neutrophil superoxide generation ( $A_{2a}$ ) and mast cell degranulation ( $A_{2b}$  and  $A_3$ ) Therapeutic  
25 applications of antagonists include allergic and non allergic inflammation, e.g., release of histamine and other inflammatory mediators.

Gastrointestinal effects include inhibition of acid secretion  
30 ( $A_1$ ) therapeutic application may include reflux and ulcerative conditions Gastrointestinal effects also include colonic, intestinal and diarrheal disease, e.g., diarrheal disease associated with intestinal inflammation ( $A_{2b}$ ).

35 Eye disorders include retinal and optic nerve head injury and trauma related disorders ( $A_3$ ). In a preferred embodiment, the eye disorder is glaucoma.

09738645-4000

Other therapeutic applications of the compounds of the invention include treatment of obesity (lipolytic properties), hypertension, treatment of depression, sedative, anxiolytic, as antileptics and as laxatives, e.g., effecting  
5 motility without causing diarrhea.

The term "disease state" is intended to include those conditions caused by or associated with unwanted levels of adenosine, adenylyl cyclase activity, increased physiological  
10 activity associated with aberrant stimulation of adenosine receptors and/or an increase in cAMP. In one embodiment, the disease state is, for example, asthma, chronic obstructive pulmonary disease, allergic rhinitis, bronchitis, renal disorders, gastrointestinal disorders, or eye disorders.  
15 Additional examples include chronic bronchitis and cystic fibrosis. Suitable examples of inflammatory diseases include non-lymphocytic leukemia, myocardial ischaemia, angina, infarction, cerebrovascular ischaemia, intermittent claudication, critical limb ischemia, venous hypertension,  
20 varicose veins, venous ulceration and arteriosclerosis. Impaired reperfusion states include, for example, any post-surgical trauma, such as reconstructive surgery, thrombolysis or angioplasty.


25 This invention also provides a combination therapy for glycoma, comprising the compound of claims 1, 6, or 18, and a prostaglandin agonist,  $\beta$ 2-2 agonist, or a muniscrinic antagonist.

30 The language "treatment of a N-6 substituted 7-deazapurine responsive state" or "treating a N-6 substituted 7-deazapurine responsive state" is intended to include changes in a disease state or condition, as described above, such  
35 that physiological symptoms in a mammal can be significantly diminished or minimized. The language also includes control, prevention or inhibition of physiological symptoms or effects



associated with an aberrant amount of adenosine. In one preferred embodiment, the control of the disease state or condition is such that the disease state or condition is eradicated. In another preferred embodiment, the control is selective such that aberrant levels of adenosine receptor activity are controlled while other physiologic systems and parameters are unaffected.

The term "N-6 substituted 7-deazapurine" is art recognized and is intended to include those compounds having the formula I:



(I)

"N-substituted 7-deazapurine" includes pharmaceutically acceptable salts thereof, and, in one embodiment, also includes certain N-6 substituted purines described herein.

In certain embodiments, the N-6 substituted 7-deazapurine is not N-6 benzyl or N-6 phenylethyl substituted. In other embodiments, R<sub>4</sub> is not benzyl or phenylethyl substituted. In preferred embodiments, R<sub>1</sub> and R<sub>2</sub> are both not hydrogen atoms. In still other preferred embodiments, R<sub>3</sub> is not a hydrogen atom.

00738546-1000

The language "therapeutically effective amount" of an N-6 substituted 7-deazapurine, described *infra*, is that amount of a therapeutic compound necessary or sufficient to perform its intended function within a mammal, e.g., treat a N-6 substituted 7-deazapurine responsive state, or a disease state in a mammal. An effective amount of the therapeutic compound can vary according to factors such as the amount of the causative agent already present in the mammal, the age, sex, and weight of the mammal, and the ability of the therapeutic compounds of the present invention to affect a N-6 substituted 7-deazapurine responsive state in the mammal. One of ordinary skill in the art would be able to study the aforementioned factors and make a determination regarding the effective amount of the therapeutic compound without undue experimentation. An *in vitro* or *in vivo* assay also can be used to determine an "effective amount" of the therapeutic compounds described *infra*. The ordinarily skilled artisan would select an appropriate amount of the therapeutic compound for use in the aforementioned assay or as a therapeutic treatment.

A therapeutically effective amount preferably diminishes at least one symptom or effect associated with the N-6 substituted 7-deazapurine responsive state or condition being treated by at least about 20%, (more preferably by at least about 40%, even more preferably by at least about 60%, and still more preferably by at least about 80%) relative to untreated subjects. Assays can be designed by one skilled in the art to measure the diminishment of such symptoms and/or effects. Any art recognized assay capable of measuring such parameters are intended to be included as part of this invention. For example, if asthma is the state being treated, then the volume of air expended from the lungs of a subject can be measured before and after treatment for measurement of increase in the volume using an art recognized technique. Likewise, if inflammation is the state being

treated, then the area which is inflamed can be measured before and after treatment for measurement of diminishment in the area inflamed using an art recognized technique.

- 5 The term "cell" includes both prokaryotic and eukaryotic cells.

The term "animal" includes any organism with adenosine receptors or any organism susceptible to a N-6-substituted 7-  
10 deazapurine responsive state. Examples of animals include yeast, mammals, reptiles, and birds. It also includes transgenic animals.

The term "mammal" is art recognized and is intended to  
15 include an animal, more preferably a warm-blooded animal, most preferably cattle, sheep, pigs, horses, dogs, cats, rats, mice, and humans. Mammals susceptible to a N-6 substituted 7-deazapurine responsive state, inflammation, emphysema, asthma, central nervous system conditions, or  
20 acute respiratory distress syndrome, for example, are included as part of this invention.

In another aspect, the present invention pertains to methods for modulating an adenosine receptor(s) in a mammal by  
25 administering to the mammal a therapeutically effective amount of a N-6 substituted 7-deazapurine, such that modulation of the adenosine receptor in the mammal occurs. Suitable adenosine receptors include the families of A<sub>1</sub>, A<sub>2</sub>, or A<sub>3</sub>. In a preferred embodiment, the N-6 substituted 7-  
30 deazapurine is an adenosine receptor antagonist.

The language "modulating an adenosine receptor" is intended to include those instances where a compound interacts with an adenosine receptor(s), causing increased, decreased or  
35 abnormal physiological activity associated with an adenosine receptor or subsequent cascade effects resulting from the modulation of the adenosine receptor. Physiological activities associated with adenosine receptors include


induction of sedation, vasodilation, suppression of cardiac rate and contractility, inhibition of platelet aggregability, stimulation of gluconeogenesis, inhibition of lipolysis, opening of potassium channels, reducing flux of calcium channels, etc.

The terms "modulate", "modulating" and "modulation" are intended to include preventing, eradicating, or inhibiting the resulting increase of undesired physiological activity associated with abnormal stimulation of an adenosine receptor, e.g., in the context of the therapeutic methods of the invention. In another embodiment, the term modulate includes antagonistic effects, e.g., diminishment of the activity or production of mediators of allergy and allergic inflammation which results from the overstimulation of adenosine receptor(s). For example, the therapeutic deazapurines of the invention can interact with an adenosine receptor to inhibit, for example, adenylate cyclase activity.

The language "condition characterized by aberrant adenosine receptor activity" is intended to include those diseases, disorders or conditions which are associated with aberrant stimulation of an adenosine receptor, in that the stimulation of the receptor causes a biochemical and or physiological chain of events that is directly or indirectly associated with the disease, disorder or condition. This stimulation of an adenosine receptor does not have to be the sole causative agent of the disease, disorder or condition but merely be responsible for causing some of the symptoms typically associated with the disease, disorder, or condition being treated. The aberrant stimulation of the receptor can be the sole factor or at least one other agent can be involved in the state being treated. Examples of conditions include those disease states listed *supra*, including inflammation, gastrointestinal disorders and those symptoms manifested by the presence of increased adenosine receptor activity. Preferred examples include those symptoms associated with

asthma, allergic rhinitis, chronic obstructive pulmonary disease, emphysema, bronchitis, gastrointestinal disorders and glaucoma.

5 The language "treating or treatment of a condition characterized by aberrant adenosine receptor activity" is intended to include the alleviation of or diminishment of at least one symptom typically associated with the condition. The treatment also includes alleviation or diminishment of  
10 more than one symptom. Preferably, the treatment cures, e.g., substantially eliminates, the symptoms associated with the condition. The present invention pertains to compounds, N-6 substituted 7-deazapurines, having the formula I:



(I)

25 wherein R<sub>1</sub> and R<sub>2</sub> are each independently a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety or together form a substituted or unsubstituted heterocyclic ring; R<sub>3</sub> is a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or  
30 alkylaryl moiety; R<sub>4</sub> is a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety. R<sub>5</sub> and R<sub>6</sub> are each independently a halogen atom, e.g., chlorine, fluorine, or bromine, a hydrogen atom or a substituted or unsubstituted alkyl, aryl, or alkylaryl  
35 moiety or R<sub>4</sub> and R<sub>5</sub> or R<sub>5</sub> and R<sub>6</sub> together form a substituted or unsubstituted heterocyclic or carbocyclic ring. Also included, are pharmaceutically acceptable

salts of the N-6 substituted 7-deazapurines.

In certain embodiments,  $R_1$  and  $R_2$  can each independently be a substituted or unsubstituted cycloalkyl or heteroarylalkyl  
5 moieties. In other embodiments,  $R_3$  is a hydrogen atom or a substituted or unsubstituted heteroaryl moiety. In still other embodiments,  $R_4$ ,  $R_5$  and  $R_6$  can each be independently a heteroaryl moiety.

10 In one embodiment,  $R_1$  is a hydrogen atom,  $R_2$  is a substituted or unsubstituted cyclohexane, cyclopentyl, cyclobutyl or cyclopropane moiety,  $R_3$  is a substituted or unsubstituted phenyl moiety,  $R_4$  is a hydrogen atom and  $R_5$  and  $R_6$  are both methyl groups.

15 In another embodiment,  $R_2$  is a cyclohexanol, a cyclohexanediol, a cyclohexylsulfonamide, a cyclohexanamide, a cyclohexylester, a cyclohexene, a cyclopentanol or a cyclopentanediol and  $R_3$  is a phenyl moiety.

20 In still another embodiment,  $R_1$  is a hydrogen atom,  $R_2$  is a cyclohexanol,  $R_3$  is a substituted or unsubstituted phenyl, pyridine, furan, cyclopentane, or thiophene moiety,  $R_4$  is a hydrogen atom, a substituted alkyl, aryl or arylalkyl moiety,  
25 and  $R_5$  and  $R_6$  are each independently a hydrogen atom, or a substituted or unsubstituted alkyl, aryl, or alkylaryl moiety.


In yet another embodiment,  $R_1$  is a hydrogen atom,  $R_2$  is  
30 substituted or unsubstituted alkylamine, arylamine, or alkylarylamine, a substituted or unsubstituted alkylamide, arylamide or alkylarylamine, a substituted or unsubstituted alkylsulfonamide, arylsulfonamide or alkylarylsulfonamide, a substituted or unsubstituted alkylurea, arylurea or  
35 alkylarylurea, a substituted or unsubstituted alkylcarbamate, arylcarbamate or alkylarylcarbamate, a substituted or unsubstituted alkylcarboxylic acid, arylcarboxylic acid or alkylarylcarboxylic acid,  $R_3$  is a substituted or unsubstituted

0072354-1990  
0072354-1990

phenyl moiety,  $R_4$  is a hydrogen atom and  $R_5$  and  $R_6$  are methyl groups.

In still another embodiment,  $R_2$  is guanidine, a modified  
5 guanidine, cyanoguanidine, a thiourea, a thioamide or an  
amidine.

In one embodiment,  $R_2$  can be



10

wherein  $R_{2a}$ - $R_{2c}$  are each independently a hydrogen atom or a  
saturated or unsaturated alkyl, aryl or alkylaryl moiety and  
 $R_{2d}$  is a hydrogen atom or a saturated or unsaturated alkyl,  
15 aryl, or alkylaryl moiety,  $NR_{2e}R_{2f}$ , or  $OR_{2g}$ , wherein  $R_{2e}$ - $R_{2g}$   
are each independently a hydrogen atom or a saturated or  
unsaturated alkyl, aryl or alkylaryl moieties.  
Alternatively,  $R_{2a}$  and  $R_{2b}$  together can form a carbocyclic or  
heterocyclic ring having a ring size between about 3 and 8  
20 members, e.g., cyclopropyl, cyclopentyl, cyclohexyl groups.

In one aspect of the invention, both  $R_5$  and  $R_6$  are not methyl  
groups, preferably, one of  $R_5$  and  $R_6$  is an alkyl group, e.g.,  
a methyl group, and the other is a hydrogen atom.


25

In another aspect of the invention, when  $R_4$  is 1-phenylethyl  
and  $R_1$  is a hydrogen atom, then  $R_3$  is not phenyl, 2-  
chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-  
dichlorophenyl, 3-methoxyphenyl or 4-methoxyphenyl or when  $R_4$   
30 and  $R_1$  are 1-phenylethyl, then  $R_3$  is not a hydrogen atom or  
when  $R_4$  is a hydrogen atom and  $R_3$  is a phenyl, then  $R_1$  is not  
phenylethyl.

35

In another aspect of the invention,  
when  $R_5$  and  $R_6$  together form a  
carbocyclic ring, e.g.,


5



or pyrimido[4,5-6]indole, then  $R_3$  is not phenyl when  $R_4$  is 1-  
10 (4-methylphenyl)ethyl, phenylisopropyl, phenyl or 1-  
phenylethyl or when  $R_3$  is not a hydrogen atom when  $R_4$  is 1-  
phenylethyl. The carbocyclic ring formed by  $R_5$  and  $R_6$  can be  
either aromatic or aliphatic and can have between 4 and 12  
carbon atoms, e.g., naphthyl, phenylcyclohexyl, etc.,  
15 preferably between 5 and 7 carbon atoms, e.g., cyclopentyl or  
cyclohexyl. Alternatively,  $R_5$  and  $R_6$  together can form a  
heterocyclic ring, such as those disclosed below. Typical  
heterocyclic rings include between 4 and 12 carbon atoms,  
preferably between 5 and 7 carbon atoms, and can be either  
20 aromatic or aliphatic. The heterocyclic ring can be further  
substituted, including substitution of one or more carbon  
atoms of the ring structure with one or more heteroatoms.

In still another aspect of the invention,  $R_1$  and  $R_2$  form a  
25 heterocyclic ring. Representative examples include, but are  
not limited to, those heterocyclic rings listed below, such  
as morpholino, piperazine and the like, e.g., 4-  
hydroxypiperidines, 4-aminopiperidines. Where  $R_1$  and  $R_2$   
together form a piperazino group,

30




35



wherein  $R_7$  can be a hydrogen atom or a substituted or unsubstituted alkyl, aryl or alkylaryl moiety.

- 5 In yet another aspect of the invention  $R_4$  and  $R_5$  together can form a heterocyclic ring, e.g.,

10



- 15 wherein the heterocyclic ring can be either aromatic or aliphatic and can form a ring having between 4 and 12 carbon atoms, e.g., naphthyl, phenylcyclohexyl, etc. and can be either aromatic or aliphatic, e.g., cyclohexyl, cyclopentyl. The heterocyclic ring can be further substituted, including substitution of carbon atoms of the ring structure with one or more heteroatoms. Alternatively,  $R_4$  and  $R_5$  together can form a heterocyclic ring, such as those disclosed below.

- 20 In certain embodiments, the N-6 substituted 7-deazapurine is not N-6 benzyl or N-6 phenylethyl substituted. In other embodiments,  $R_4$  is not benzyl or phenylethyl substituted. In preferred embodiments,  $R_1$  and  $R_2$  are both not hydrogen atoms. In still other preferred embodiments,  $R_3$  is not H.

- 30 The compounds of the invention may comprise water-soluble prodrugs which are described in WO 99/33815, International Application No. PCT/US98/04595, filed March 9, 1998 and published July 8, 1999. The entire content of WO 99/33815 is expressly incorporated herein by reference. The water-soluble prodrugs are metabolized in vivo to an active drug, e.g., by esterase catalyzed hydrolysis. Examples of potential prodrugs include deazapurines with, for example,  $R_2$  as cycloalkyl substituted with  $-OC(O)(Z)NH_2$ , wherein Z is a
- 35

side chain of a naturally or unnaturally occurring amino acid, or analog thereof, an  $\alpha$ ,  $\beta$ ,  $\gamma$ , or  $\omega$  amino acids, or a dipeptide. Preferred amino acid side chains include those of glycine, alanine, valine, leucine, isoleucine, lysine,  $\alpha$ -methylalanine, aminocyclopropane carboxylic acid, azetidine-2-carboxylic acid,  $\beta$ -alanine,  $\gamma$ -aminobutyric acid, alanine-alanine, or glycine-alanine.

In a further embodiment, the invention features deazapurines of the formula (I), wherein  $R_1$  is hydrogen;  $R_2$  is substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkyl, or  $R_1$  and  $R_2$  together form a substituted or unsubstituted heterocyclic ring,  $R_3$  is unsubstituted or substituted aryl;  $R_4$  is hydrogen; and  $R_5$  and  $R_6$  are each independently hydrogen or alkyl, and pharmaceutically acceptable salts thereof. The deazapurines of this embodiment may potentially be selective  $A_3$  receptor antagonists.

In one embodiment,  $R_2$  is substituted (e.g., hydroxy substituted) or unsubstituted cycloalkyl. In an advantageous subembodiment,  $R_1$  and  $R_4$  are hydrogen,  $R_3$  is unsubstituted or substituted phenyl, and  $R_5$  and  $R_6$  are each alkyl. Preferably  $R_2$  is mono-hydroxycyclopentyl or mono-hydroxycyclohexyl.  $R_2$  also may be substituted with  $-\text{NH}-\text{C}(=\text{O})\text{E}$ , wherein E is substituted or unsubstituted  $\text{C}_1$ - $\text{C}_4$  alkyl (e.g., alkylamine, e.g., ethylamine.).

$R_1$  and  $R_2$  may also together form a substituted or unsubstituted heterocyclic ring, which may be substituted with an amine or acetamido group.

In another aspect,  $R_2$  may be  $-\text{A}-\text{NHC}(=\text{O})\text{B}$ , wherein A is unsubstituted  $\text{C}_1$ - $\text{C}_4$  alkyl (e.g., ethyl, propyl, butyl), and B is substituted or unsubstituted  $\text{C}_1$ - $\text{C}_4$  alkyl (e.g., methyl, aminoalkyl, e.g., aminomethyl or aminoethyl, alkylamino, e.g., methylamino, ethylamino), preferably when  $R_1$  and  $R_4$  are

hydrogen,  $R_3$  is unsubstituted or substituted phenyl, and  $R_5$  and  $R_6$  are each alkyl. B may be substituted or unsubstituted cycloalkyl, e.g., cyclopropyl or 1-amino-cyclopropyl.

5 In another embodiment,  $R_3$  may be substituted or unsubstituted phenyl, preferably when  $R_5$  and  $R_6$  are each alkyl. Preferably,  $R_3$  may have one or more substituents (e.g., *o*-, *m*- or *p*-chlorophenyl, *o*-, *m*- or *p*-fluorophenyl).

10 Advantageously,  $R_3$  may be substituted or unsubstituted heteroaryl, preferably when  $R_5$  and  $R_6$  are each alkyl. Examples of heteroaryl groups include pyridyl, pyrimidyl, pyridazinyl, pyrazinyl, pyrrolyl, triazolyl, thioazolyl, oxazolyl, oxadiazolyl, furanyl, methylenedioxyphenyl and  
15 thiophenyl. Preferably,  $R_3$  is 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl or 3-pyrimidyl.

Preferably in one embodiment,  $R_5$  and  $R_6$  are each hydrogen. In another,  $R_5$  and  $R_6$  are each methyl.

20 In a particularly preferred embodiment, the deazapurines of the invention are water-soluble prodrugs that can be metabolized *in vivo* to an active drug, e.g. by esterase catalyzed hydrolysis. Preferably the prodrug comprises an  $R_2$   
25 group which is cycloalkyl substituted with  $-OC(O)(Z)NH_2$ , wherein Z is a side chain of a naturally or unnaturally occurring amino acid, an analog thereof, an  $\alpha$ ,  $\beta$ ,  $\gamma$ , or  $\omega$  amino acid, or a dipeptide. Examples of preferred side chains include the side chains of glycine, alanine, valine,  
30 leucine, isoleucine, lysine,  $\alpha$ -methylalanine, aminocyclopropane carboxylic acid, azetidine-2-carboxylic acid,  $\beta$ -alanine,  $\gamma$ -aminobutyric acid, alanine-alanine, or glycine-alanine.

35 In a particularly preferred embodiment, Z is a side chain of glycine,  $R_2$  is cyclohexyl,  $R_3$  is phenyl, and  $R_5$  and  $R_6$  are methyl.

00729545-120400  
007027-932760

In another embodiment, the deazapurine is 4-(*cis*-3-hydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine.

- 5 In another embodiment, the deazapurine is 4-(*cis*-3-(2-aminoacetoxy) cyclopentyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine trifluoroacetic acid salt.

10 In another embodiment, the deazapurine is 4-(3-acetamido)piperidiny-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine.

15 In another embodiment, the deazapurine is 4-(2-*N*'-methylureapropyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine.

20 In another embodiment, the deazapurine is 4-(2-acetamidobutyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine.

In another embodiment, the deazapurine is 4-(2-*N*'-methylureabutyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine.

25 In another embodiment, the deazapurine is 4-(2-aminocyclopropylacetamidoethyl)amino-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine.

30 In another embodiment, the deazapurine is 4-(*trans*-4-hydroxycyclohexyl)amino-2-(3-chlorophenyl)-7*H*-pyrrolo[2,3*d*] pyrimidine.

35 In another embodiment, the deazapurine is 4-(*trans*-4-hydroxycyclohexyl)amino-2-(3-fluorophenyl)-7*H*-pyrrolo[2,3*d*] pyrimidine.

09733645-149400

In another embodiment, the deazapurine is 4-(trans-4-hydroxycyclohexyl)amino-2-(4-pyridyl)-7H-pyrrolo[2,3d]pyrimidine.

- 5 In yet another embodiment, the invention features a method for inhibiting the activity of an adenosine receptor (e.g., A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub>, or, preferably, A<sub>3</sub>) in a cell, by contacting the cell with N-6 substituted 7-deazapurine (e.g., preferably, an adenosine receptor antagonist).
- 10 In another aspect, the invention features a method for treating damage to the eye of an animal (e.g., a human) by administering to the animal an effective amount of an N-6 substituted 7-deazapurine. Preferably, the N-6 substituted
- 15 7-deazapurine is an antagonist of A<sub>3</sub> adenosine receptors in cells of the animal. The damage is to the retina or the optic nerve head and may be acute or chronic. The damage may be the result of, for example, glaucoma, edema, ischemia, hypoxia or trauma.
- 20 In a preferred embodiment, the invention features a deazapurine having the formula II, supra, wherein X is N or CR<sub>6</sub>; R<sub>1</sub> and R<sub>2</sub> are each independently hydrogen, or substituted or unsubstituted alkoxy, aminoalkyl, alkyl, aryl, or
- 25 alkylaryl, or together form a substituted or unsubstituted heterocyclic ring, provided that both R<sub>1</sub> and R<sub>2</sub> are both not hydrogen; R<sub>3</sub> is substituted or unsubstituted alkyl, arylalkyl, or aryl; R<sub>4</sub> is hydrogen or substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl; L is hydrogen, substituted or unsubstituted alkyl, or
- 30 R<sub>4</sub> and L together form a substituted or unsubstituted heterocyclic or carbocyclic ring; R<sub>6</sub> is hydrogen, substituted or unsubstituted alkyl, or halogen; Q is CH<sub>2</sub>, O, S, or NR<sub>7</sub>, wherein R<sub>7</sub> is hydrogen or substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl; and W is unsubstituted or substituted alkyl,
- 35 cycloalkyl, alkynyl, aryl, arylalkyl, biaryl, heteroaryl, substituted carbonyl, substituted thiocarbonyl, or substituted sulfonyl, provided that if R<sub>3</sub> is pyrrolidino, then

007866 46 490400

R<sub>4</sub> is not methyl.

In one embodiment, in compounds of formula II, X is CR, and Q is CH<sub>2</sub>, O, S, or NH. In another embodiment, X is N.

5

In a further embodiment of compounds of formula II, W is substituted or unsubstituted aryl, 5- or 6- member heteroaryl, or biaryl. W may be substituted with one or more substituents. Examples of substituents include: halogen, hydroxy, alkoxy, amino, aminoalkyl, aminocarboxamide, CN, CF<sub>3</sub>, CO<sub>2</sub>R<sub>8</sub>, CONHR<sub>8</sub>, CONR<sub>8</sub>R<sub>9</sub>, SOR<sub>8</sub>, SO<sub>2</sub>R<sub>8</sub>, and SO<sub>2</sub>NR<sub>8</sub>R<sub>9</sub>, wherein R<sub>8</sub> and R<sub>9</sub> are each independently hydrogen, or substituted or unsubstituted alkyl, cycloalkyl, aryl, or arylalkyl. Preferably, W may be substituted or unsubstituted phenyl, e.g., methylenedioxyphenyl. W also may be a substituted or unsubstituted 5-membered heteroaryl ring, e.g., pyrrole, pyrazole, oxazole, imidazole, triazole, tetrazole, furan, thiophene, thiazole, and oxadiazole. Preferably, W may be a 6-member heteroaryl ring, e.g., pyridyl, pyrimidyl, pyridazinyl, pyrazinal, and thiophenyl. In a preferred embodiment, W is 2-pyridyl, 3- pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, or 5-pyrimidyl.

In one advantageous embodiment of compounds of formula II, Q is NH and W is a 3-pyrazolo ring which is unsubstituted or N-substituted by substituted or unsubstituted alkyl, cycloalkyl, aryl, or arylalkyl.

In another embodiment of compounds of formula II, Q is oxygen, and W is a 2-thiazolo ring which is unsubstituted or substituted by substituted or unsubstituted alkyl, cycloalkyl, aryl, or arylalkyl.

In another embodiment of compounds of formula II, W is substituted or unsubstituted alkyl, cycloalkyl e.g., cyclopentyl, or arylalkyl. Examples of substituents include halogen, hydroxy, substituted or unsubstituted alkyl,




alkoxy, substituted or unsubstituted alkylamine, arylamine,  
or alkylarylamine, substituted or unsubstituted aminoalkyl,  
amino aryl, or aminoalkylaryl, substituted or unsubstituted  
alkylamide, arylamide or alkylarylamide, substituted or  
5 unsubstituted alkylsulfonamide, arylsulfonamide or  
alkylarylsulfonamide, substituted or unsubstituted alkylurea,  
arylurea or alkylarylurea, substituted or unsubstituted  
alkylcarbamate, arylcarbamate or alkylarylcabamate, or  
substituted or unsubstituted alkylcarboxylic acid,  
10 arylcarboxylic acid or alkylarylcaboxylic acid.

Preferably,  $R_2$  is substituted or unsubstituted cycloalkyl,  
e.g., mono- or dihydroxy-substituted cyclohexyl or  
cyclopentyl (preferably, monohydroxy-substituted cyclohexyl  
15 or monohydroxy-substituted cyclopentyl).


DocId: 32326660



Advantageously,  $R_2$  may be of the following formula:



wherein A is  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_7$  cycloalkyl, a chain of one to seven atoms, or a ring of three to seven atoms, optionally substituted with  $C_1$ - $C_6$  alkyl, halogens, hydroxyl, carboxyl, thiol, or amino groups; wherein B is methyl,  $N(Me)_2$ ,  $N(Et)_2$ , NHMe, NHEt,  $(CH_2)_rNH_3^+$ ,  $NH(CH_2)_rCH_3$ ,  $(CH_2)_rNH_2$ ,  $(CH_2)_rCHCH_3NH_2$ ,  $(CH_2)_rNHMe$ ,  $(CH_2)_rOH$ ,  $CH_2CN$ ,  $(CH_2)_mCO_2H$ ,  $CHR_{18}R_{19}$ , or  $CHMeOH$ , wherein r is an integer from 0 to 2, m is 1 or 2,  $R_{18}$  is alkyl,  $R_{19}$  is  $NH_3^+$  or  $CO_2H$  or  $R_{18}$  and  $R_{19}$  together are:



wherein p is 2 or 3; and  $R_{17}$  is  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_7$  cycloalkyl, a chain of one to seven atoms, or a ring of three to seven atoms, optionally substituted with  $C_1$ - $C_6$  alkyl, halogens, hydroxyl, carboxyl, thiol, or amino groups.


Advantageously, A is unsubstituted or substituted  $C_1$ - $C_6$  alkyl. B may be unsubstituted or unsubstituted  $C_1$ - $C_6$  alkyl.

In a preferred embodiment,  $R_2$  is of the formula  $-A-NHC(=O)B$ . In a particularly advantageous embodiment, A is  $-CH_2CH_2-$  and B is methyl.

The compounds of the invention may comprise water-soluble prodrugs which are metabolized *in vivo* to an active drug, e.g., by esterase catalyzed hydrolysis. Examples of potential prodrugs include deazapurines with, for example,  $R_2$  as cycloalkyl substituted with  $-OC(O)(Z)NH_2$ , wherein Z is a

side chain of a naturally or unnaturally occurring amino acid, or analog thereof, an  $\alpha$ ,  $\beta$ ,  $\gamma$ , or  $\omega$  amino acid, or a dipeptide. Preferred amino acid side chains include those of glycine, alanine, valine, leucine, isoleucine, lysine,  $\alpha$ -methylalanine, aminocyclopropane carboxylic acid, azetidine-2-carboxylic acid,  $\beta$ -alanine,  $\gamma$ -aminobutyric acid, alanine-alanine, or glycine-alanine.

In another embodiment,  $R_1$  and  $R_2$  together are:



wherein  $n$  is 1 or 2, and wherein the ring may be optionally substituted with one or more hydroxyl, amino, thiol, carboxyl, halogen,  $CH_2OH$ ,  $CH_2NHC(=O)alkyl$ , or  $CH_2NHC(=O)NHalkyl$  groups. Preferably,  $n$  is 1 or 2 and said ring is substituted with  $-NHC(=O)alkyl$ .

In one advantageous embodiment,  $R_1$  is hydrogen,  $R_2$  is substituted or unsubstituted  $C_1-C_6$  alkyl,  $R_3$  is substituted or unsubstituted phenyl,  $R_4$  is hydrogen,  $L$  is hydrogen or substituted or unsubstituted  $C_1-C_6$  alkyl,  $Q$  is O, S or  $NR_7$ , wherein  $R_7$  is hydrogen or substituted or unsubstituted  $C_1-C_6$  alkyl, and  $W$  is substituted or unsubstituted aryl. Preferably,  $R_2$  is  $-A-NHC(=O)B$ , wherein  $A$  and  $B$  are each independently unsubstituted or substituted  $C_1-C_4$  alkyl. For example,  $A$  may be  $CH_2CH_2$ .  $B$  may be, for example, alkyl (e.g., methyl), or aminoalkyl (e.g., aminomethyl). Preferably,  $R_3$  is unsubstituted phenyl and  $L$  is hydrogen.  $R_6$  may be methyl or preferably, hydrogen. Preferably,  $Q$  is O, S, or  $NR_7$  wherein  $R_7$  is hydrogen or substituted or unsubstituted  $C_1-C_6$  alkyl, e.g., methyl.  $W$  is unsubstituted or substituted phenyl (e.g., alkoxy, halogen substituted). Preferably,  $W$  is

p-fluorophenyl, p-chlorophenyl, or p-methoxyphenyl. W may also be heteroaryl, e.g., 2-pyridyl.

In a particularly preferred embodiment, the deazapurine is 4-  
5 (2-acetylaminoethyl) amino-6-phenoxyethyl-2-phenyl-7H-  
pyrrolo[2,3d]pyrimidine.

In a particularly preferred embodiment, the deazapurine is 4-  
(2-acetylaminoethyl) amino-6-(4-fluorophenoxy)methyl-2-  
10 phenyl-7H-pyrrolo[2,3d]pyrimidine.

In a particularly preferred embodiment, the deazapurine is  
4-(2-acetylaminoethyl) amino-6-(4-chlorophenoxy)methyl-2-  
phenyl-7H-pyrrolo[2,3d]pyrimidine.

15 In a particularly preferred embodiment, the deazapurine is 4-  
(2-acetylaminoethyl) amino-6-(4-methoxyphenoxy)methyl-2-  
phenyl-7H-pyrrolo[2,3d]pyrimidine.

20 In a particularly preferred embodiment, the deazapurine is 4-  
(2-acetylaminoethyl) amino-6-(2-pyridyloxy)methyl-2-phenyl-  
7H-pyrrolo[2,3d]pyrimidine.

In a particularly preferred embodiment, the deazapurine is 4-  
25 (2-acetylaminoethyl) amino-6-(N-phenylamino)methyl-2-phenyl-  
7H-pyrrolo[2,3d]pyrimidine.

In a particularly preferred embodiment, the deazapurine is 4-  
(2-acetylaminoethyl) amino-6-(N-methyl-N-phenylamino)methyl-  
30 2-phenyl-7H-pyrrolo[2,3d]pyrimidine.

In a particularly preferred embodiment, the deazapurine is 4-  
(2-N'-methylureaethyl) amino-6-phenoxyethyl-2-phenyl-7H-  
pyrrolo[2,3d]pyrimidine.

35 The invention further pertains to a method for inhibiting the

DocId: 3433460

activity of an adenosine receptor (e.g., an  $A_{2r}$  adenosine receptor) in a cell by contacting the cell with a compound of the invention. Preferably, the compound is an antagonist of the receptor.

5

The invention also pertains to a method for treating a gastrointestinal disorder (e.g., diarrhea) in an animal by administering to an animal an effective amount of a compound of the invention (e.g., an antagonist of  $A_{2r}$ ). Preferably,

10 the animal is a human.

In another embodiment, the invention relates to a pharmaceutical composition containing an N-6 substituted 7-deazapurine of the invention and a pharmaceutically

15 acceptable carrier.

The invention also pertains to a method for treating a N-6 substituted 7-deazapurine responsive state in an animal, by administering to a mammal a therapeutically effective amount of a deazapurine of the invention, such that treatment of a N-6 substituted 7-deazapurine responsive state in the animal occurs. Advantageously, the disease state may be a disorder mediated by adenosine. Examples of preferred disease states include: central nervous system disorders, cardiovascular disorders, renal disorders, inflammatory disorders, allergic disorders, gastrointestinal disorders, eye disorders, and respiratory disorders.

The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. The term alkyl further includes alkyl groups, which can further include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone, e.g., oxygen, nitrogen, sulfur or phosphorous atoms. In preferred embodiments, a straight

chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., C<sub>1</sub>-C<sub>30</sub> for straight chain, C<sub>3</sub>-C<sub>30</sub> for branched chain), and more preferably 20 or fewer. Likewise, preferred cycloalkyls have from 4-10 carbon atoms in their ring structure, and more preferably have 5, 6 or 7 carbons in the ring structure.

Moreover, the term alkyl as used throughout the specification and claims is intended to include both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate. Cycloalkyls can be further substituted, e.g., with the substituents described above. An "alkylaryl" moiety is an alkyl substituted with an aryl (e.g., phenylmethyl (benzyl)). The term "alkyl" also includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

The term "aryl" as used herein, refers to the radical of aryl groups, including 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms, for

example, benzene, pyrrole, furan, thiophene, imidazole, benzoxazole, benzothiazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Aryl groups also include polycyclic fused aromatic groups such as naphthyl, quinolyl, indolyl, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heteroaryls" or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinate, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with alicyclic or heterocyclic rings which are not aromatic so as to form a polycycle (e.g., tetralin).

The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively. For example, the invention contemplates cyano and propargyl groups.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, more preferably from one to six carbon atoms in its backbone structure, even more preferably one to three carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths.

The terms "alkoxyalkyl", "polyaminoalkyl" and "thioalkoxyalkyl" refer to alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms replacing one or more carbons of the hydrocarbon backbone, 5 e.g., oxygen, nitrogen or sulfur atoms.

The terms "polycyclyl" or "polycyclic radical" refer to the radical of two or more cyclic rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in 10 which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, 15 hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, 20 dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, 25 heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms 30 are nitrogen, oxygen, sulfur and phosphorus.

The term "amino acids" includes naturally and unnaturally occurring amino acids found in proteins such as glycine, alanine, valine, cysteine, leucine, isoleucine, serine, 35 threonine, methionine, glutamic acid, aspartic acid, glutamine, asparagine, lysine, arginine, proline, histidine, phenylalanine, tyrosine, and tryptophan. Amino acid analogs

007334 40400

include amino acids with lengthened or shortened side chains or variant side chains with appropriate functional groups. Amino acids also include D and L stereoisomers of an amino acid when the structure of the amino acid admits of stereoisomeric forms. The term "dipeptide" includes two or more amino acids linked together. Preferably, dipeptides are two amino acids linked via a peptide linkage. Particularly preferred dipeptides include, for example, alanine-alanine and glycine-alanine.

10

It will be noted that the structure of some of the compounds of this invention includes asymmetric carbon atoms and thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. All such isomeric forms of these compounds are expressly included in this invention. Each stereogenic carbon may be of the R or S configuration. It is to be understood accordingly that the isomers arising from such asymmetry (e.g., all enantiomers and diastereomers) are included within the scope of this invention, unless indicated otherwise. Such isomers can be obtained in substantially pure form by classical separation techniques and by stereochemically controlled synthesis.

25 The invention further pertains to pharmaceutical compositions for treating a N-6 substituted 7-deazapurine responsive state in a mammal, e.g., respiratory disorders (e.g., asthma, bronchitis, chronic obstructive pulmonary disorder, and allergic rhinitis), renal disorders, gastrointestinal disorders, and eye disorders. The pharmaceutical composition includes a therapeutically effective amount of a N-6 substituted 7-deazapurine, described supra, and a pharmaceutically acceptable carrier. It is to be understood, that all of the deazapurines described above are included for therapeutic treatment. It is to be further understood that the deazapurines of the invention can be used alone or in combination with other deazapurines of the invention or in



combination with additional therapeutic compounds, such as antibiotics, antiinflammatories, or anticancer agents, for example.

5 The term "antibiotic" is art recognized and is intended to include those substances produced by growing microorganisms and synthetic derivatives thereof, which eliminate or inhibit growth of pathogens and are selectively toxic to the pathogen while producing minimal or no deleterious effects upon the  
10 infected host subject. Suitable examples of antibiotics include, but are not limited to, the principle classes of aminoglycosides, cephalosporins, chloramphenicols, fuscidic acids, macrolides, penicillins, polymyxins, tetracyclines and streptomycins.

15 The term "antiinflammatory" is art recognized and is intended to include those agents which act on body mechanisms, without directly antagonizing the causative agent of the inflammation such as glucocorticoids, aspirin, ibuprofen, NSAIDS, etc.

20 The term "anticancer agent" is art recognized and is intended to include those agents which diminish, eradicate, or prevent growth of cancer cells without, preferably, adversely affecting other physiological functions. Representative  
25 examples include cisplatin and cyclophosphamide.

When the compounds of the present invention are administered as pharmaceuticals, to humans and mammals, they can be given per se or as a pharmaceutical composition containing, for  
30 example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

The phrase "pharmaceutically acceptable carrier" as used  
35 herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a compound(s) of the

00703646-120400

As set out above, certain embodiments of the present compounds can contain a basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable acids. The term "pharmaceutically acceptable salts" in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate,

acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and  
5 the like. (See, e.g., Berge et al. (1977) "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19).

In other cases, the compounds of the present invention may contain one or more acidic functional groups and, thus, are  
10 capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable bases. The term "pharmaceutically acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These  
15 salts can likewise be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation,  
20 with ammonia, or with a pharmaceutically acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation  
25 of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

The term "pharmaceutically acceptable esters" refers to the  
30 relatively non-toxic, esterified products of the compounds of the present invention. These esters can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form or hydroxyl with a suitable esterifying agent.  
35 Carboxylic acids can be converted into esters via treatment with an alcohol in the presence of a catalyst. Hydroxyl containing derivatives can be converted into esters via

treatment with an esterifying agent such as alkanoyl halides. The term is further intended to include lower hydrocarbon groups capable of being solvated under physiological conditions, e.g., alkyl esters, methyl, ethyl and propyl  
5 esters. (See, for example, Berge et al., *supra*.)

The invention further contemplates the use of prodrugs which are converted *in vivo* to the therapeutic compounds of the invention (see, e.g., R.B. Silverman, 1992, "The Organic  
10 Chemistry of Drug Design and Drug Action", Academic Press, Chapter 8). Such prodrugs can be used to alter the biodistribution (e.g., to allow compounds which would not typically enter the reactive site of the protease) or the pharmacokinetics of the therapeutic compound. For example,  
15 a carboxylic acid group, can be esterified, e.g., with a methyl group or an ethyl group to yield an ester. When the ester is administered to a subject, the ester is cleaved, enzymatically or non-enzymatically, reductively or hydrolytically, to reveal the anionic group. An anionic  
20 group can be esterified with moieties (e.g., acyloxymethyl esters) which are cleaved to reveal an intermediate compound which subsequently decomposes to yield the active compound. In another embodiment, the prodrug is a reduced form of a sulfate or sulfonate, e.g., a thiol, which is oxidized in  
25 *vivo* to the therapeutic compound. Furthermore, an anionic moiety can be esterified to a group which is actively transported *in vivo*, or which is selectively taken up by target organs. The ester can be selected to allow specific targeting of the therapeutic moieties to particular reactive  
30 sites, as described below for carrier moieties.

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring  
35 and perfuming agents, preservatives and antioxidants can also be present in the compositions.

00729516-1 200400

5

15

20

35

liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing  
5 a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; solution retarding agents, such as paraffin; absorption accelerators, such as quaternary ammonium compounds; wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; absorbents, such as kaolin and bentonite clay; lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

35 A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent,

preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert dilutents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl

alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert dilutents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be



The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the active compound in a polymer matrix or gel.

The ophthalmic formulations of the present invention may include one or more deazapurines and a pharmaceutically acceptable vehicle. Various types of vehicles may be used. The vehicles will generally be aqueous in nature. Aqueous solutions are generally preferred, based on ease of

formulation, as well as a patient's ability to easily administer such compositions by means of instilling one to two drops of the solutions in the affected eyes. However, the deazapurines of the present invention may also be readily  
5 incorporated into other types of compositions, such as suspensions, viscous or semi-viscous gels or other types of solid or semi-solid compositions. The ophthalmic compositions of the present invention may also include various other ingredients, such as buffers, preservatives,  
10 co-solvents and viscosity building agents.

An appropriate buffer system (e.g., sodium phosphate, sodium acetate or sodium borate) may be added to prevent pH drift under storage conditions.

15 Ophthalmic products are typically packaged in multidose form. Preservatives are thus required to prevent microbial contamination during use. Suitable preservatives include: benzalkonium chloride, thimerosal, chlorobutanol, methyl  
20 paraben, propyl paraben, phenylethyl alcohol, edetate disodium, sorbic acid, polyquaternium-1, or other agents known to those skilled in the art. Such preservatives are typically employed at a level of from 0.001 to 1.0% weight/volume ("% w/v").

25 When the deazapurines of the present invention are administered during intraocular surgical procedures, such as through retrobulbar or periocular injection and intraocular perfusion or injection, the use of balanced salt irrigating  
30 solutions as vehicles are most preferred. BSS® Sterile Irrigating Solution and BSS Plus® Sterile Intraocular Irrigating Solution (Alcon Laboratories, Inc., Fort Worth, Texas, USA) are examples of physiologically balanced intraocular irrigating solutions. The latter type of  
35 solution is described in U.S. Pat. No. 4,550,022 (Garabedian, et al.), the entire contents of which are hereby incorporated in the present specification by reference. Retrobulbar and periocular injections are known to those skilled in the art

00723646-4040  
00723646-4040

and are described in numerous publications including, for example, *Ophthalmic Surgery: Principles of Practice*, Ed., G. L. Spaeth. W. B. Sanders Co., Philadelphia, Pa., U.S.A., pages 85-87 (1990).

5

As indicated above, use of deazapurines to prevent or reduce damage to retinal and optic nerve head tissues at the cellular level is a particularly important aspect of one embodiment of the invention. Ophthalmic conditions which may be treated include, but are not limited to, retinopathies, macular degeneration, ocular ischemia, glaucoma, and damage associated with injuries to ophthalmic tissues, such as ischemia reperfusion injuries, photochemical injuries, and injuries associated with ocular surgery, particularly injuries to the retina or optic nerve head by exposure to light or surgical instruments. The compounds may also be used as an adjunct to ophthalmic surgery, such as by vitreal or subconjunctival injection following ophthalmic surgery. The compounds may be used for acute treatment of temporary conditions, or may be administered chronically, especially in the case of degenerative disease. The compounds may also be used prophylactically, especially prior to ocular surgery or noninvasive ophthalmic procedures, or other types of surgery.

25 Pharmaceutical compositions of this invention suitable for parenteral administration comprise one or more compounds of the invention in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

35

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol,

03729545 44400  
00729545 44400

5

10

20

35

5 The preparations of the present invention may be given orally, parenterally, topically, or rectally. They are of course given by forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, 10 suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories. Oral administration is preferred.

The phrases "systemic administration," "administered systematically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

These compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray, 35 rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

35 In general, a suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such

an effective dose will generally depend upon the factors described above. Generally, intravenous and subcutaneous doses of the compounds of this invention for a patient, when used for the indicated analgesic effects, will range from  
5 about 0.0001 to about 200 mg per kilogram of body weight per day, more preferably from about 0.01 to about 150 mg per kg per day, and still more preferably from about 0.2 to about 140 mg per kg per day.

10 If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms.

15 While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical composition.

The present invention also pertains to packaged  
20 pharmaceutical compositions for treating a N-6 substituted 7 deazapurine responsive state, e.g., undesirable increased adenosine receptor activity in a mammal. The packaged pharmaceutical compositions include a container holding a therapeutically effective amount of at least one deazapurine  
25 as described *supra* and instructions for using the deazapurine for treating the deazapurine responsive state in the mammal.

The deazapurines of the invention can be prepared using standard methods for organic synthesis. Deazapurines can be  
30 purified by reverse phase HPLC, chromatography, recrystallization, etc. and their structures confirmed by mass spectral analysis, elemental analysis, IR and/or NMR spectroscopy.

35 Typically, synthesis of the intermediates as well as the deazapurines of the invention is performed in solution. The addition and removal of one or more protecting group is also


00728616-420400

typical practice and is known to those skilled in the art. Typical synthetic schemes for the preparation of deazapurine intermediates of the invention are outlined below in Scheme I.

5


This invention also provides a method of preparing compound IV, comprising the steps of

10




wherein P is a removable protecting group;


b) treating the product of step a) under cyclization conditions to provide



c) treating the product of step b) under suitable conditions to provide



d) treating the chlorinated product of step c) with  $\text{NHR}_1\text{R}_2$  to provide






wherein R<sub>1</sub> is H and R<sub>2</sub> is cyclopropyl methylamino  
carbonylethyl, *cis*-3-hydroxy cyclopentyl, acetamido  
butyl, methylamino carbonylamino butyl, ethylamino  
carbonylamino propyl, methylamino carbonylamino propyl,  
5 2-acetyl amino-3-methyl butyl, N,N-diethylamino  
carbonylamino ethyl, thioacetamido ethyl, 3-amino  
acetyloxy cyclopentyl, 3-hydroxy cyclopentyl, 2-pyrrolyl  
carbonyl aminoethyl, 2-imidazolidinone ethyl, 1-  
aminocarbonyl-2-methyl propyl, 1-aminocarbonyl-2-phenyl  
10 ethyl, 3-hydroxy azetidino, 2-imidazolyl ethyl,  
acetamido ethyl, 1-(R)-phenyl-2-hydroxyethyl, N-  
methylaminocarbonyl pyridyl-2-methyl, or R<sub>1</sub>, R<sub>2</sub> and the  
nitrogen together are 3-acetamido piperadino, 3-hydroxy  
pyrrolidino, 3-methyloxy carbonylmethyl pyrrolidino, 3-  
15 aminocarbonylmethyl pyrrolidino, or 3-hydroxymethyl  
piperadino.

wherein R<sub>3</sub> is a substituted or unsubstituted four to six  
membered ring;

20 wherein R<sub>4</sub> is H, alkyl, substituted alkyl, or cycloalkyl;


wherein R<sub>5</sub> is H, alkyl, substituted alkyl, aryl, or  
substituted aryl.

25 This invention also provides a method of preparing compound  
of V, comprising the steps of




wherein P is a removable protecting group;


b) treating the product of step a) under cyclization conditions to provide



c) treating the product of step b) under suitable conditions to provide



d) treating the chlorinated product of step c) with  $\text{NH}_2\text{CH}_2(\text{CH}_2)_m\text{CH}_2\text{NHC}(=\text{O})\text{R}_1$  to provide



wherein m is 0, 1, or 2;

5 wherein R<sub>1</sub> is cyclopropyl methyl, methyl, methylamino, or  
aminomethyl;

wherein R<sub>2</sub> is aryl, substituted aryl, heteroaryl;


10 wherein R<sub>3</sub> is H, alkyl, substituted alkyl, or cycloalkyl;

15 wherein R<sub>6</sub> is H, alkyl, substituted alkyl, aryl,  
arylalkyl, amino, substituted aryl, wherein said  
substituted alkyl is -C(R<sub>9</sub>)(R<sub>10</sub>)NR<sub>7</sub>R<sub>8</sub>, wherein R<sub>9</sub> and R<sub>10</sub>  
are each H or alkyl, wherein R<sub>7</sub> and R<sub>8</sub> are each alkyl or  
cycloalkyl, or R<sub>7</sub>, R<sub>8</sub> and the nitrogen together form a  
ring system of between 4 and 7 members.


20 This invention further provided a method of preparing  
compound VI, comprising

0079556-4000  
0079556-4000

a) reacting




to provide




wherein P is a removable protecting group;

b) treating the product of step a) under cyclization conditions to provide




c) treating the product of step b) under suitable conditions to provide




; and

d) treating the chlorinated product of step c) with



to provide




wherein  $R_2$  is unsubstituted aryl.

wherein  $R_3$  is H, alkyl, substituted alkyl, or cycloalkyl;

5 wherein  $R_4$  H, alkyl, substituted alkyl, aryl, arylalkyl, amino, substituted aryl, wherein said substituted alkyl is  $-C(R_9)(R_{10})NR_7R_8$ , wherein  $R_9$  and  $R_{10}$  are each H or alkyl, wherein  $R_7$  and  $R_8$  are each alkyl or cycloalkyl, or  $R_7$ ,  $R_8$  and the nitrogen together form a ring system of  
10 between 4 and 7 members.

This invention also provides a compound having the structure:



IV

wherein  $R_1$  is H and  $R_2$  is cyclopropyl methylamino  
25 carbonylethyl, *cis*-3-hydroxy cyclopentyl, acetamido  
butyl, methylamino carbonylamino butyl, ethylamino  
carbonylamino propyl, methylamino carbonylamino propyl,  
2-acetyl amino-3-methyl butyl, N,N-diethylamino  
carbonylamino ethyl, thioacetamido ethyl, 3-amino  
30 acetyloxy cyclopentyl, 3-hydroxy cyclopentyl; 2-pyrrolyl  
carbonyl aminoethyl, 2-imidazolidinone ethyl, 1-  
aminocarbonyl-2-methyl propyl, 1-aminocarbonyl-2-phenyl  
ethyl, 3-hydroxy azetidino, 2-imidazolyl ethyl,  
acetamido ethyl, 1-(R)-phenyl-2-hydroxyethyl, N-  
35 methylaminocarbonyl pyridyl-2- methyl, or  $R_1$ ,  $R_2$  and the  
nitrogen together are 3-acetamido piperadino, 3-hydroxy  
pyrrolidino, 3-methyloxy carbonylmethyl pyrrolidino, 3-  
aminocarbonylmethyl pyrrolidino, or 3-hydroxymethyl


piperadino.

wherein R<sub>3</sub> is a substituted or unsubstituted benzene,  
pyrrole, thiophene, furan, thiazole, imidazole,  
5 pyrazole, 1,2,4-triazole, pyridine, 2(1H)-pyridone,  
4(1H)-pyridone, pyrazine, pyrimidine, pyridazine,  
isothiazole, isoxazole, oxazole, tetrazole, naphthalene,  
tetralin, naphthyridine, benzofuran, benzothiophene,  
indole, 2,3-dihydroindole, 1H-indole, indoline,  
10 benzopyrazole, 1,3-benzodioxole, benzoxazole, purine,  
coumarin, chromone, quinoline, tetrahydroquinoline,  
isoquinoline, benzimidazole, quinazoline, pyrido[2,3-  
b]pyrazine, pyrido[3,4-b]pyrazine, pyrido[3,2-  
c]pyridazine, purido[3,4-b]-pyridine, 1H-pyrazole[3,4-  
15 d]pyrimidine, pteridine, 2(1H)-quinolone, 1(2H)-  
isoquinolone, 1,4-benzisoxazine, benzothiazole,  
quinoxaline, quinoline-N-oxide, isoquinoline-N-oxide,  
quinoxaline-N-oxide, quinazoline-N-oxide, benzoxazine,  
phthalazine, or cinnoline.

wherein R<sub>5</sub> is H, alkyl, substituted alkyl, or cycloalkyl;

wherein R<sub>6</sub> is H, alkyl, substituted alkyl, aryl, or  
substituted aryl.

In one embodiment of the compound, the compound has the  
structure:




In another embodiment of the compound, R<sub>3</sub> is phenyl.

In another embodiment of the compound, R<sub>3</sub> is hydrogen or methyl.

5

In another embodiment of the compound, R<sub>3</sub> is hydrogen, methyl, phenyl, 3-chlorophenyloxy methyl, or trans-2- phenylamino methyl pyrrolidino methyl.

10 This invention further provides a compound having the structure:



V

25

wherein m is 0, 1, or 2;

wherein R<sub>1</sub> is cyclopropyl methyl, methyl, methylamino, or aminomethyl;

30

wherein R<sub>2</sub> is aryl, substituted aryl, or heteroaryl;

wherein R<sub>5</sub> is H, alkyl, substituted alkyl, or cycloalkyl;

35

wherein R<sub>6</sub> is H, alkyl, substituted alkyl, aryl, arylalkyl, amino, substituted aryl, wherein said substituted alkyl is -C(R<sub>9</sub>)(R<sub>10</sub>)NR<sub>7</sub>R<sub>8</sub>, wherein R<sub>9</sub> and R<sub>10</sub> are each H or alkyl, wherein R<sub>7</sub> and R<sub>8</sub> are each alkyl or

cycloalkyl, or R<sup>-</sup>, R<sub>2</sub> and the nitrogen together form a ring system of between 4 and 7 members.

In one embodiment of compound V, m is 0 and R<sub>2</sub> is phenyl.

5

In another embodiment of compound V, m is 1 and R<sub>2</sub> is phenyl.

In another embodiment of compound V, m is 2 and R<sub>2</sub> is phenyl.

10 In another embodiment of compound V, R<sub>5</sub> and R<sub>6</sub> are methyl.

In another embodiment of compound V, R<sub>5</sub> and R<sub>6</sub> are methyl.

In another embodiment of compound V, R<sub>5</sub> and R<sub>6</sub> are methyl.


15

In another embodiment of compound V, the compound has the structure:

20

25

30




35

(Compound 1316)




In another embodiment of compound V, the compound has the structure:




(Compound 1311)

In another embodiment of compound V, the compound has the structure:




(Compound 1202)

In another embodiment of compound V, the compound has the structure:




(Compound 1310)

In another embodiment of compound V, the compound has the structure:




(Compound 1312)

This invention further provides a compound having the structure:



(Compound 609)

This invention also provides a compound having the structure:




VI

wherein  $R_2$  is unsubstituted aryl.

wherein  $R_5$  is H, alkyl, substituted alkyl, or cycloalkyl;


wherein  $R_6$  H, alkyl, substituted alkyl, aryl, arylalkyl, amino, substituted aryl, wherein said substituted alkyl is  $-C(R_9)(R_{10})NR_7R_8$ , wherein  $R_9$  and  $R_{10}$  are each H or alkyl, wherein  $R_7$  and  $R_8$  are each alkyl or cycloalkyl, or  $R_7$ ,  $R_8$  and the nitrogen together form a ring system of between 4 and 7 members.

25 In one embodiment of compound VI, the compound has the structure:




(Compound 1309)

In one embodiment of compound 1309, the compound has the structure:




In another embodiment of compound 1309, the compound has the structure:



35

This invention also provides a compound having the structure:




VII

wherein R<sub>1</sub> is 3-hydroxy cyclopentyl ethylamino carbonylamino propyl, N,N-diethylamino carbonylamino ethyl, thioacetamido ethyl, 3-amino acetyloxy cyclopentyl, 3-hydroxy cyclopentyl, 2-pyrrolyl carbonyl aminoethyl, 2-imidazolidinone ethyl, 1-aminocarbonyl-2-methyl propyl, 1-aminocarbonyl-2-phenyl ethyl, 3-hydroxy azetidino, 2-imidazolyl ethyl, acetamido ethyl, 1-(R)-phenyl-2-hydroxyethyl, or N-methylaminocarbonyl pyridyl-2-methyl;


wherein R<sub>3</sub> and R<sub>4</sub> are independently H, substituted or unsubstituted alkyl, or aryl.

In one embodiment of the compound, the compound has the structure:




(Compound 1700)

In another embodiment of the compound, the compound has the structure:




(Compound 1701)

In another embodiment of the compound, the compound has the structure:



(Compound 1702)

In another embodiment of the compound, the compound has the structure:



(Compound 1704)




In another embodiment of the compound, the compound has the structure:

5

10

15



(Compound 1705)


20

In another embodiment of the compound, the compound has the structure:

25


30

35




(Compound 1706)


In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:




25 In another embodiment of the compound, the compound has the structure:



In another embodiment of the compound, the compound has the structure:




15 In another embodiment of the compound, the compound has the structure:



(Compound 1707)


00739646 180400

In another embodiment of the compound, the compound has the structure:




(Compound 1708)

In another embodiment of the compound, the compound has the structure:



(Compound 1709)


In another embodiment of the compound, the compound has the structure:



15

(Compound 1710)

In another embodiment of the compound, the compound has the structure:




(Compound 1712)

35

In another embodiment of the compound, the compound has the structure:


5




(Compound 1713)

20 In another embodiment of the compound, the compound has the structure:

25




In another embodiment of the compound, the compound has the structure:




(Compound 1714)

In another embodiment of the compound, the compound has the structure:




35

In another embodiment of the compound, the compound has the structure:




(Compound 1715)

15 In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:





This invention also provides a compound having the structure:



VIII

15


wherein R<sub>1</sub>, R<sub>2</sub> and the nitrogen together are 3-hydroxy pyrrolidino, 3-methoxy carbonylmethyl pyrrolidino, 3-aminocarbonylmethyl pyrrolidino, or 3-hydroxymethyl piperadino;

20

wherein R<sub>2</sub> and R<sub>4</sub> are independently H, substituted or unsubstituted alkyl, or aryl.


In one embodiment of the compound, the compound has the

25 structure:




(Compound 1711)

In another embodiment of the compound, the compound has the structure:




15 (Compound 1703)

In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:




007345-40400

In another embodiment of the compound, the compound has the structure:




(Compound 1716)


In another embodiment of the compound, the compound has the structure:



In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:




35

(Compound 1717)


In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:



In another embodiment of the compound, the compound has the structure:




In another embodiment of the compound, the compound has the structure:



35

In another embodiment of the compound, the compound has the structure:



This invention also provides a method for treating a disease associated with A<sub>3</sub> adenosine receptor in a subject, comprising administering to the subject a therapeutically effective amount of any of the compounds IV, V, VI, VI, or VIII.

In one embodiment of the method, the subject is a mammal.

In another embodiment of the method, the mammal is a human.

In another embodiment of the method, said A<sub>3</sub> adenosine receptor is associated with a central nervous system disorder, a cardiovascular disorder, asthma, hypersensitivity, rhinitis, hay fever, serum sickness, allergic vasculitis, atopic dermantitis, dermantitis, psoriasis, eczema, idiopathic pulmonary fibrosis, eosinophilic chloreycystitis, chronic airway inflammation, hypereosinophilic syndromes, eosinophilic gastroenteritis, edema, urticaria, eosinophilic myocardial disease, episodic angioedema with eosinophilia, inflammatory bowel disease, ulcerative colitis, allergic granulomatosis, carcinomatosis,



eosinophilic granuloma, familial histiocytosis, hypertension, mast cell degranulation, tumor, cardiac hypoxia, cerebral ischemia, diuresis, renal failure, neurological disorder, mental disorder, cognitive disorder, myocardial ischemia, 5 bronchoconstriction, arthritis, autoimmune disease, Crohn's disease, Grave's disease, diabetes, multiple sclerosis, anaemia, psoriasis, fertility disorders, lupus erythematosus, reperfusion injury, brain arteriole diameter, the release of allergic mediators, scleroderma, stroke, global ischemia, 10 central nervous system disorder, cardiovascular disorder, renal disorder, inflammatory disorder, gastrointestinal disorder, eye disorder, allergic disorder, respiratory disorder, or immunological disorder.

15 Diseases associated with adenosine A<sub>1</sub>, A<sub>2a</sub>, A<sub>2b</sub> and A<sub>3</sub> receptors are disclosed in WO 99/06053 and WO-09822465, WO-09705138, WO-09511681, WO-09733879, JP-09291089, PCT/US98/16053 and U.S. Patent No. 5,516,894, the entire content of which are fully incorporate herein by reference.

20 This invention also provides a water-soluble prodrug of any of the compounds IV, V, VI, VII, or VIII; wherein said water-soluble prodrug that is metabolized *in vivo* to an active drug which selectively inhibit A<sub>3</sub> adenosine receptor.

25 In one embodiment of the prodrug, said prodrug is metabolized *in vivo* by esterase catalyzed hydrolysis.

This invention also provides a pharmaceutical composition 30 comprising the prodrug and a pharmaceutically acceptable carrier.

This invention also provides a method for inhibiting the activity of an A<sub>3</sub> adenosine receptor in a cell, which 35 comprises contacting said cell with any of the compounds IV, V, VI, VII, or VIII.

09334400

In one embodiment of the method, the compound is an antagonist of said A<sub>3</sub> adenosine receptor.

In another embodiment of the pharmaceutical composition, said  
5 pharmaceutical composition is an ophthalmic formulation.

In another embodiment of the pharmaceutical composition, said pharmaceutical composition is an periocular, retrobulbar or intraocular injection formulation.

10

In another embodiment of the pharmaceutical composition, said pharmaceutical composition is a systemic formulation.

This invention also provides a method for treating a  
15 gastrointestinal disorder in an subject, comprising administering to the an effective amount of any of the compounds IV, V, VI, VII, or VIII.

In one embodiment of the method, said disorder is diarrhea.  
20

In another embodiment of the method, the subject is a human.

In another embodiment of the method, the compound is an antagonist of A<sub>3</sub> adenosine receptors.

25

This invention further provides a method for treating respiratory disorder in a subject, comprising administering to the subject an effective amount of any of the compounds IV, V, VI, VII, or VIII.

30

In one embodiment of the method, said disorder is asthma, chronic obstructive pulmonary disease, allergic rhinitis, or an upper respiratory disorder.

35 In another embodiment of the method, the subject is a human.

In another embodiment of the method, said compound is an

antagonist of A<sub>3</sub> adenosine receptors.

This invention also provides a method for treating damage to the eye of a subject which comprises administering to said  
5 subject an effective amount of any of the compounds IV, V, VI, VII, or VIII.

In one embodiment of the method, said damage comprises retinal or optic nerve head damage.  
10

In another embodiment of the method, said damage is acute or chronic.

In another embodiment of the method, said damage is the  
15 result of glaucoma, edema, ischemia, hypoxia or trauma.

In another embodiment of the method, the subject is a human.

In another embodiment of the method, the compound is an  
20 antagonist of A<sub>3</sub> adenosine receptors.

This invention also provide a pharmaceutical composition comprising a therapeutically effective amount of any of the compounds IV, V, VI, VII, or VIII and a pharmaceutically  
25 acceptable carrier.

In one embodiment of the pharmaceutical composition, said therapeutically effective amount is effective to treat a respiratory disorder or a gastrointestinal disorder.  
30

In another embodiment of the pharmaceutical composition, said gastrointestinal disorder is diarrhea.

In another embodiment of the pharmaceutical composition,  
35 said respiratory disorder is asthma, allergic rhinitis, or chronic obstructive pulmonary disease.

In another embodiment of the pharmaceutical composition,

said pharmaceutical composition is an ophthalmic formulation.

In another embodiment of the pharmaceutical composition, said pharmaceutical composition is an periocular, retrobulbar or  
5 intraocular injection formulation.

In another embodiment of the pharmaceutical composition, said pharmaceutical composition is a systemic formulation.

10 In another embodiment of the pharmaceutical composition, said pharmaceutical composition is a surgical irrigating solution.

This inventio also provides a packaged pharmaceutical  
15 composition for treating a disease associated with A<sub>3</sub> adenosine receptor in a subject, comprising: (a) a container holding a therapeutically effective amount of any of the compounds IV, V, VI, VII, or VIII; and (b) instructions for using said compound for treating said disease in a subject.

20 Compounds represented by the formula IV, V, VI, VII, and VIII can be synthesized by the Schemes I-IX.

As used herein, "A compound is A<sub>3</sub> selective." means that a  
25 compound has a binding constant to adenosine A<sub>3</sub> receptor of at least ten time higher then that to adenosine A<sub>1</sub>, A<sub>2a</sub>, or A<sub>2b</sub>.

The invention is further illustrated by the following  
30 examples which in no way should be construed as being further limiting. The contents of all references, pending patent applications and published patent applications, cited throughout this application, including those referenced in the background section, are hereby incorporated by reference.

35 It should be understood that the models used throughout the examples are accepted models and that the demonstration of efficacy in these models is predictive of efficacy in humans.

00729516-120400




# EXPERIMENTAL DETAILS

The deazapurines of the invention can be prepared using standard methods for organic synthesis. Deazapurines can be purified by reverse phase HPLC, chromatography, recrystallization, etc. and their structures confirmed by mass spectral analysis, elemental analysis, IR and/or NMR spectroscopy.

Typically, synthesis of the intermediates as well as the deazapurines of the invention is performed in solution. The addition and removal of one or more protecting group is also typical practice and is known to those skilled in the art. Typical synthetic schemes for the preparation of deazapurine intermediates of the invention are outlined below in Scheme I.

**Scheme I**



wherein  $R_3$ ,  $R_5$  and  $R_6$  are as defined above.















In general, a protected 2-amino-3-cyano-pyrrole can be treated with an acyl halide to form a carboxyamido-3-cyano-pyrrole which can be treated with acidic methanol to effect ring closure to a pyrrolo[2,3d]pyrimidine-4(3H)-one (Muller, C.E. et al. J. Med. Chem. 40:4396 (1997)). Removal of the pyrrolo protecting group followed by treatment with a chlorinating reagent, e.g., phosphorous oxychloride, produced substituted or unsubstituted 4-chloro-7H-pyrrolo[2,3d]pyrimidines. Treatment of the chloropyrimidine with amines afforded 7-deazapurines.












For example, as shown in Scheme I, a N-(1-dl-phenylethyl)-2-amino-3-cyano-pyrrole was treated with an acyl halide in pyridine and dichloromethane. The resultant N-(1-dl-phenylethyl)-2-phenylcarboxyamido-3-cyano-pyrrole was treated with a 10:1 mixture of methanol/sulfuric acid to effect ring closure, resulting in a dl-7H-7-(1-phenylethyl)pyrrolo[2,3d]pyrimidine-4(3H)-one. Removal of the phenylethyl group by treatment of the pyrimidine with polyphosphoric acid (PPA) followed by  $POCl_3$  afforded a key intermediate, the 4-chloro-7H-pyrrolo[2,3d]pyrimidine. Further treatment of the 4-chloro-7H-pyrrolo[2,3d]pyrimidine with various amines listed in Table 1 gives compounds of formula (I) and (II).

TABLE 1

R	M <sup>+</sup> + H	R	M <sup>+</sup> + H
	343.2		351.27
	343.18		430.35
	337.21		359.44
	364.19		404.32
	330.18		330.45
	347.22		339.47
	350.28		353.41




	344.19		324.45
	394.16		359.38
	371.12		379.40
	359.39		387.41
	403.33		344.48
	351.49		337.53
	330.37		295.2

	407.23		321.2
	355.45		337.53
	441.33		350.2
	413.24		343.2
	372.48		373.2
			307.2

A general approach to prepare 6-substituted pyrroles is depicted in the following scheme (Scheme II).

Scheme II




wherein  $R_1$  through  $R_5$  are as defined above.

Transesterification and alkylation of ethyl cyanoacetate with  
 35 an  $\alpha$ -haloketone affords a ketomethylester. Protection of the  
 ketone followed by treatment with an amidine (e.g., alkyl,  
 aryl or alkylaryl) hydrochloride produced the resultant ketal  
 protected pyrimidine. Removal of the protecting group,  
 followed by cyclization and treatment with phosphorous  
 40 oxychloride afforded the chloride intermediate which could be  
 further treated with an amine to afford an amine 6-

substituted pyrrole. Additionally, alkylation of the pyrrole nitrogen can be achieved under art recognized conditions.

A general approach to prepare 5-substituted pyrroles is depicted in the following scheme (Scheme III).

Scheme III




wherein R<sub>1</sub> through R<sub>6</sub> are defined as above and R is a removable protecting group.

Condensation of malononitrile and an excess of a ketone followed by bromination of the product afforded a mixture of starting material, monobrominated and dibrominated products which were treated with an alkylamine, arylamine or alkylarylamine. The resultant amine product was acylated with an acid chloride and the monacylated pyrrole was

cyclized in the presence of acid to afford the corresponding pyrimidine. The pyrrole protecting group was removed with polyphosphoric acid and treated with phosphorous oxychloride to produce a chlorinated product. The chlorinated pyrrole could subsequently be treated with an amine to produce an amino 5-substituted pyrrole. Alkylation of the pyrrole nitrogen can be achieved under art recognized conditions.

Schemes IV and V depict methods for preparing the deazapurines 1 and 2 of the invention.




wherein  $R_5$  and  $R_6$  are as described above, e.g.,  $CH_3$ .

## 2) Specific Preparation of 6-methyl pyrrolopyrimidines:

The key reaction toward 6-methylpyrrolopyrimidines (1) [ $R_5 = CH_3$ ] was cyclization of a cyanoacetate with benzamidine to a pyrimidine. It was believed methyl cyanoacetate would cyclize more efficiently with benzamidine to a pyrimidine than the corresponding ethyl ester. Therefore, transesterification and alkylation of ethyl cyanoacetate in the presence of NaOMe and an excess of an  $\alpha$ -haloacetyl moiety, e.g., chloroacetone, gave the desired methyl ester (3) in 79% yield (Scheme IV). The ketoester (3) was protected as the acetal (4) in 81% yield. A new cyclization method to the pyrimidine (5) was achieved with an amidine hydrochloride, e.g., benzamidine hydrochloride, with 2

equivalents of DBU to afford the 5 in 54% isolated yield. This method improves the yield from 20% using the published conditions, which utilizes NaOMe during the cyclization with guanidine. Cyclization to the pyrrole-pyrimidine (6) was achieved via deprotection of the acetal in aqueous HCl in 78% yield. Reaction of (6) with phosphorous oxychloride at reflux gave the corresponding 4-chloro derivative (7). Coupling with *trans*-4-aminocyclohexanol in dimethyl sulfoxide at 135°C gave (1) in 57% from (7). One skilled in the art will appreciate that choice of reagents allows for great flexibility in choosing the desired substituent R<sub>5</sub>.

Scheme IV



Specific Preparation of 5-methylpyrrolopyrimidines

Knöevenagel condensation of malononitrile and an excess ketone, e.g., acetone in refluxing benzene gave 8 in 50% yield after distillation. Bromination of 8 with N-bromosuccinimide in the presence of benzoyl peroxide in chloroform yielded a mixture of starting material, mono- (9), and di-brominated products (5/90/5) after distillation (70%). The mixture was reacted with an  $\alpha$ -methylalkylamine or  $\alpha$ -methylarylamine, e.g.,  $\alpha$ -methylbenzylamine, to deliver the aminopyrrole (10). After passing through a short silica gel column, the partially purified amine (31% yield) was acylated with an acid chloride, e.g., benzoyl chloride to deliver mono- (11), and diacylated (12) pyrroles, which were separated by flash chromatography. Acid hydrolysis of the disubstituted pyrrole (12) generated a combined yield of 29% for the acylpyrrole (11). Cyclization in the presence of concentrated sulphuric acid and DMF yielded (13) (23%), which was deprotected with polyphosphoric acid to (14). Reaction of (14) with phosphorous oxychloride at reflux gave the corresponding 4-chloro derivative (15). Coupling with trans-4-aminocyclohexanol in dimethyl sulfoxide at 135°C gave (2) [ $R_6 = CH_3$ ] in 30% from (14) (See Scheme V). One skilled in the art will appreciate that choice of reagents allows for great flexibility in choosing the desired substituent  $R_6$ .


25

30

35

0078846-100400

Scheme V





Alternative Synthetic Route to R<sub>6</sub>-Substituted Pyrroles, e.g.,

5-methyl pyrrolopyrimidines:

This alternative route to R<sub>6</sub>-substituted pyrroles, e.g., 5-methylpyrrolopyrimidines, involves transesterification and  
5 alkylation of ethyl cyanoacetate to (16) (Scheme VI). The condensation of (16) with benzamidine hydrochloride with 2 equivalents of DBU affords the pyrimidine (17). Cyclization to the pyrrole-pyrimidine (14) will be achieved via deprotection of the acetal in aqueous HCl. Reaction of (14)  
10 with phosphorous oxychloride at reflux gave the corresponding 4-chloro derivative (15). Coupling with trans-4-aminocyclohexanol in dimethyl sulfoxide at 135°C gives 2. This procedure reduces the number of synthetic reactions to the target compound (2) from 9 to 4 steps. Moreover, the  
15 yield is dramatically improved. Again, one skilled in the art will appreciate that choice of reagents allows for great flexibility in choosing the desired substituent R<sub>6</sub>.

20


25

30

35


0078846-10400

Scheme VI




A general approach to prepare  
des-methyl pyrrole is depicted  
in the following scheme

30 (Scheme VII)



Scheme VII




wherein  $R_1$  through  $R_3$  are defined as above.

Alkylation of an alkyl cyanoacetate with a diethyl acetal in the presence of a base afforded a cyano diethyl acetal which was treated with an amidine salt to produce a methyl pyrrolopyrimidine precursor. The precursor was chlorinated and treated with an amine to form the *des*-methyl pyrrolopyrimidine target as shown above.

For example, Scheme VIII depicts the synthesis of compound (18).

Scheme VIII








Commercially available methyl cyanoacetate was alkylated with  
 bromoacetaldehyde diethyl acetal in the presence of potassium  
 35 carbonate and NaI to yield (19). Cyclization to the  
 pyrimidine (20) was achieved in two steps. Initially, the  
 pyrimidine-acetal was formed via reaction of (19) with  
 benzamidine hydrochloride with 2 equivalents of DBU. The

resultant pyrimidine-acetal was deprotected without purification with aqueous 1 N HCl and the resultant aldehyde cyclized to the pyrrolo-pyrimidine (20), which was isolated by filtration. Reaction of (20) with phosphorous oxychloride at reflux afforded the corresponding 4-chloro derivative (21). Coupling of the chloro derivative with trans-4-aminocyclohexanol in DMSO at 135°C gave compound (18) from compound (21).

10 Schemes II-VIII demonstrate that it is possible to functionalize the 5- and 6-position of the pyrrolopyrimidine ring. Through the use of different starting reagents and slight modifications of the above reaction schemes, various functional groups can be introduced at the 5- and 6-positions in formula (I) and (II). Table 2 illustrates some examples.

Table 2. Selected list of 5- and 6-substituted pyrrolopyrimidines.

20	Starting Reagent	R <sub>5</sub>	R <sub>6</sub>
		H	
		H	Substituted Ar
		H	CH <sub>2</sub> C(O)OCH <sub>3</sub>
25		C(O)OCH <sub>3</sub>	CH <sub>3</sub>
30		C(O)NHCH <sub>3</sub>	CH <sub>3</sub>



### Exemplification

#### Preparation 1:

A modification of the alkylation method of Seela and Lüpke was used.<sup>1</sup> To an ice-cooled (0°C) solution of ethyl cyanoacetate (6.58 g, 58.1 mmol) in MeOH (20 mL) was slowly added a solution of NaOMe (25% w/v; 58.1 mmol). After 10 min, chloroacetone (5 mL; 62.8 mmol) was slowly added. After 4 h, the solvent was removed. The brown oil was diluted with EtOAc (100 mL) and washed with H<sub>2</sub>O (100 mL). The organic fraction was dried, filtered, and concentrated to a brown oil (7.79 g; 79%). The oil (3) (Scheme IV) was a mixture of methyl/ethyl ester products (9/1), and was used without further purification. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ<sub>4.24</sub> (q, *J* = 7.2 Hz, OCH<sub>2</sub>), 3.91 (dd, 1H, *J* = 7.2, 7.0 Hz, CH), 3.62 (s, 3H, OCH<sub>3</sub>), 3.42 (dd, 1H, *J* = 15.0, 7.1 Hz, 1 x CH<sub>2</sub>); 3.02 (dd, 1H, *J* = 15.0, 7.0 Hz, 1 x CH<sub>2</sub>); 2.44 (s, 3H, CH<sub>3</sub>), 1.26 (t, *J* = 7.1 Hz, ester-CH<sub>3</sub>).

<sup>1</sup>Seela, F.; Lüpke, U. Chem. Ber. 1977, 110, 1462-1469.

#### 20 Preparation 2:

The procedure of Seela and Lüpke was used.<sup>1</sup> Thus, protection of the ketone (3) (Scheme IV; 5.0 g, 32.2 mmol) with ethylene glycol (4 mL, 64.4 mmol) in the presence of TsOH (100 mg) afforded (4) as an oil (Scheme IV; 5.2 g, 81.0) after flash chromatography (SiO<sub>2</sub>; 3/7 EtOAc/Hex, *R<sub>f</sub>* 0.35). Still contains ~5% ethyl ester: <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ<sub>4.24</sub> (q, *J* = 7.2 Hz, OCH<sub>2</sub>), 3.98 (s, 4H, 2 x acetal-CH<sub>2</sub>), 3.79 (s, 3H, OCH<sub>3</sub>), 3.62 (dd, 1H, *J* = 7.2, 7.0 Hz, CH), 2.48 (dd, 1H, *J* = 15.0, 7.1 Hz, 1 x CH<sub>2</sub>), 2.32 (dd, 1H, *J* = 15.0, 7.0 Hz, 1 x CH<sub>2</sub>); 1.35 (s, 3H, CH<sub>3</sub>), 1.26 (t, *J* = 7.1 Hz, ester-CH<sub>3</sub>); MS (ES): 200.1 (M<sup>+</sup>+1).

<sup>1</sup>Seela, F.; Lüpke, U. Chem. Ber. 1977, 110, 1462-1469.

**Preparation 3:**

A solution of acetal (4) (Scheme IV, 1 g, 5.02 mmol), benzamidine (786 mg, 5.02 mmol), and DBU (1.5 mL, 10.04 mmol) in dry DMF (15 mL) was heated to 85°C for 15 h. The mixture was diluted with CHCl<sub>3</sub> (30 mL) and washed with 0.5 N NaOH (10 mL) and H<sub>2</sub>O (20 mL). The organic fraction was dried, filtered and concentrated to a brown oil. Flash chromatography (SiO<sub>2</sub>; 1/9 EtOAc/CH<sub>2</sub>Cl<sub>2</sub>, R<sub>f</sub> 0.35) was attempted, but material crystallized on the column. The silica gel was washed with MeOH. Fractions containing the product (5) (Scheme IV) were concentrated and used without further purification (783 mg, 54.3%): <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 8.24 (m, 2H, Ar-H), 7.45 (m, 3H, Ar-H), 5.24 (br s, 2H, NH<sub>2</sub>), 3.98 (s, 4H, 2 x acetal-CH<sub>2</sub>), 3.60-3.15 (m, 2H, CH<sub>2</sub>), 1.38 (s, 3H, CH<sub>3</sub>); MS (ES): 288.1 (M<sup>+</sup>+1).

Preparation of compound (20) (Scheme VIII): A solution of acetal (19) (4.43 g, 20.6 mmol)<sup>1</sup>, benzamine hydrochloride (3.22 g, 20.6 mmol), and DBU (6.15 mL, 41.2 mmol) in dry DMF (20 mL) was heated to 85°C for fifteen hours. The mixture was diluted with 100mL of CHCl<sub>3</sub>, and washed with H<sub>2</sub>O (2 x 50 mL). The organic fraction was dried, filtered, and concentrated to a dark brown oil. The dark brown oil was stirred in 1N HCl (100 mL) for 2 hours at room temperature. The resulting slurry was filtered yielding the HCl salt of (20) as a tan solid (3.60 g, 70.6%); <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) 11.92 (s 1H), 8.05 (m, 2H, Ar-H), 7.45 (m, 3H, Ar-H), 7.05 (s, 1H, pyrrole-H); MS(ES): 212.1 (M<sup>+</sup>+1).

**Preparation 4:**

A solution of acetal (5) (700 mg, 2.44 mmol) in 1 N HCl (40 mL) was stirred for 2 h at RT. The resultant slurry was filtered yielding the HCl salt of 2-phenyl-6-methyl-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one as a tan solid (498 mg, 78.0%): <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 11.78 (s, 1H), 8.05 (m, 2H, Ar-H), 7.45 (m, 3H, Ar-H), 6.17 (s, 1H, pyrrole-H), 2.25 (s, 3H, CH<sub>3</sub>); MS (ES): 226.1 (M<sup>+</sup>+1).



**Preparation 5:**

A modification of the Chen et al. cyclization method was used.<sup>1</sup> To an ice-cooled (0°C) solution of bromide (9),  
5 (Scheme V; 20.0 g, 108 mmol; 90% pure) in isopropyl alcohol (60 mL) was slowly added a solution of  $\alpha$ -methylbenzylamine (12.5 mL, 97.3 mmol). The black solution was allowed to warm to RT and stir for 15 h. The mixture was diluted with EtOAc (200 mL) and washed with 0.5 N NaOH (50 mL). The organic  
10 fraction was dried, filtered, and concentrated to a black tar (19.2 g; 94%). The residue was partially purified by flash chromatography (SiO<sub>2</sub>; 4/96 MeOH/CH<sub>2</sub>Cl<sub>2</sub>, R<sub>f</sub> 0.35) to a black solid (6.38 g, 31%) as the compound *dl*-1-(1-phenylethyl)-2-amino-3-cyano-4-methylpyrrole: MS (ES): 226.1 (M<sup>+</sup>+1).

15 <sup>1</sup>Chen, Y. L.; Mansbach, R. S.; Winter, S. M.; Brooks, E.; Collins, J.; Corman, M. L.; Dunaiskis, A. R.; Faraci, W. S.; Gallaschun, R. J.; Schmidt, A.; Schulz, D. W. *J. Med. Chem.* 1997, 40, 1749-1754.

**20 Preparation 6:**

To a solution of *dl*-1-(1-phenylethyl)-2-amino-3-cyano-4,5-dimethylpyrrole<sup>1</sup> (14.9 g, 62.5 mmol) and pyridine (10.0 mL) in dichloromethane (50.0 mL) was added benzoyl chloride (9.37 g, 66.7 mmol) at 0°C. After stirring at 0°C for 1 hr, hexane  
25 (10.0 mL) was added to help precipitation of product. Solvent was removed in vacuo and the solid was recrystallized from EtOH/H<sub>2</sub>O to give 13.9 g (65%) of *dl*-1-(1-phenylethyl)-2-phenylcarbonylamino-3-cyano-4,5-dimethylpyrrole. mp 218-221°C;  
<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  1.72 (s, 3H), 1.76 (d, J = 7.3 Hz, 3H), 1.98 (s, 3H), 5.52 (q, J = 7.3 Hz, 1H), 7.14-7.54 (m, 9H), 7.68-7.72 (dd, J = 1.4 Hz, 6.9 Hz, 2H), 10.73 (s, 1H); MS (ES): 344.4 (M<sup>+</sup>+1).

<sup>1</sup> Liebig's Ann. Chem. 1986, 1485-1505.

**Preparation 6A:**

10 *d*l-1-(1-phenylethyl)-2-(2-furyl)carbonylamino-3-cyano-4,5-  
dimethylpyrrole. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.84 (d, J = 7.4 Hz,  
3H), 1.92 (s, 3H), 2.09 (s, 3H), 5.49 (q, J = 7.4 Hz, 1H),  
6.54 (dd, J = 1.8 Hz, 3.6 Hz, 1H), 7.12-7.47 (m, 7H); MS  
(ES): 334.2 (M<sup>+</sup>+1), 230.1.

dl-1-(1-phenylethyl)-2-cyclopentylcarbonylamino-3-cyano-4,5-dimethylpyrrole. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.82 (d, J = 7.4 Hz, 3H), 1.88 (s, 3H), 2.05 (s, 3H), 1.63-1.85 (m, 8H), 2.63 (m, 1H), 5.43 (q, J = 7.4 Hz, 1H), 6.52 (s, 1H), 7.05-7.20 (m, 5H); MS (ES): 336.3 (M<sup>+</sup>+1).

d1-1-(1-phenylethyl)-2-(3-thienyl) carbonylamino-3-cyano-4,5-dimethylpyrrole.

35 <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.83 (d, J = 7.0 Hz, 3H), 1.99 (s, 3H), 2.12 (s, 3H), 5.49 (q, J = 7.0 Hz, 1H), 6.90 (m, 1H), 7.18-7.36 (m, 6H), 7.79 (m, 1H); MS (ES): 350.2 (M<sup>+</sup>+1), 246.1.

dl-1-(1-phenylethyl)-2-(4-fluorophenyl)carbonylamino-3-cyano-4,5-dimethylpyrrole.

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.83 (d, J = 7.4 Hz, 3H), 1.96 (s, 3H), 2.08 (s, 3H), 5.51 (q, J = 7.4 Hz, 1H), 7.16-7.55 (m, 9H); MS (ES): 362.2 (M<sup>+</sup>+1), 258.1.

dl-1-(1-phenylethyl)-2-(3-fluorophenyl)carbonylamino-3-cyano-4,5-dimethylpyrrole.

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.83 (d, J = 7.4 Hz, 3H), 1.97 (s, 3H), 2.10 (s, 3H), 5.50 (q, J = 7.4 Hz, 1H), 7.05-7.38 (m, 7H), 7.67-7.74 (m, 2H); MS (ES): 362.2 (M<sup>+</sup>+1), 258.1.

dl-1-(1-phenylethyl)-2-(2-fluorophenyl)carbonylamino-3-cyano-4,5-dimethylpyrrole.

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.85 (d, J = 7.2 Hz, 3H), 1.94 (s, 3H), 2.11 (s, 3H), 5.50 (q, J = 7.2 Hz, 1H), 7.12-7.35 (m, 6H), 7.53 (m, 1H), 7.77 (m, 1H), 8.13 (m, 1H); MS (ES): 362.2 (M<sup>+</sup>+1), 258.0.

dl-1-(1-phenylethyl)-2-isopropylcarbonylamino-3-cyano-4,5-

dimethylpyrrole. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.19 (d, J = 7.0 Hz, 6H), 1.82 (d, J = 7.2 Hz, 3H), 1.88 (s, 3H), 2.06 (s, 3H), 2.46 (m, 1H), 5.39 (m, J = 7.2 Hz, 1H), 6.64 (s, 1H), 7.11-7.36 (m, 5H); MS (ES): 310.2 (M<sup>+</sup>+1), 206.1 .

In the case of acylation of dl-1-(1-phenylethyl)-2-amino-3-cyano-4-methylpyrrole, monoacylated dl-1-(1-phenylethyl)-2-benzoylamino-3-cyano-4-dimethylpyrrole and diacylated pyrrole dl-1-(1-phenylethyl)-2-dibenzoylamino-3-cyano-4-methylpyrrole were obtained. Monoacylated pyrrole: <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)

δ 7.69 (d, 2H, J = 7.8 Hz, Ar-H), 7.58-7.12 (m, 8H, Ar-H), 6.18 (s, 1H, pyrrole-H), 5.52 (q, 1H, J = 7.2 Hz, CH-CH<sub>3</sub>), 2.05 (s, 3H, pyrrole-CH<sub>3</sub>), 1.85 (d, 3H, J = 7.2 Hz, CH-CH<sub>3</sub>);

MS (ES): 330.2 (M<sup>+</sup>+1); Diacylated pyrrole: <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 7.85 (d, 2H, J = 7.7 Hz, Ar-H), 7.74 (d, 2H, J = 7.8

Hz, Ar-H), 7.52-7.20 (m, 9H, Ar-H), 7.04 (m, 2H, Ar-H), 6.21 (s, 1H, pyrrole-H), 5.52 (q, 1H, J = 7.2 Hz, CH-CH<sub>3</sub>), 1.77 (d,

3H,  $J = 7.2$  Hz, CH-CH<sub>3</sub>), 1.74 (s, 3H, pyrrole-CH<sub>3</sub>); MS (ES): 434.1 (M<sup>+</sup>+1).

**Preparation 7:**

- 5 To a solution of *dl*-1-(1-phenylethyl)-2-phenylcarboxyamido-3-cyano-4,5-dimethylpyrrole (1.0 g, 2.92 mmol) in methanol (10.0 mL) was added concentrated sulfuric acid (1.0 mL) at 0°C. The resulted mixture was refluxed for 15 hr and cooled down to room temperature. The precipitate was filtered to  
10 give 0.48 g (48%) of *dl*-5,6-dimethyl-2-phenyl-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 2.02 (d,  $J = 7.4$  Hz, 3H), 2.04 (s, 3H), 2.41 (s, 3H), 6.25 (q,  $J = 7.4$  Hz, 1H), 7.22-7.50 (m, 9H), 8.07-8.12 (dd,  $J = 3.4$  Hz, 6.8 Hz, 2H), 10.51 (s, 1H); MS (ES): 344.2 (M<sup>+</sup>+1).  
15 The following compounds were obtained in a similar manner as that of Preparation 7:

- dl*-5,6-dimethyl-2-(3-pyridyl)-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  
20 δ 2.03 (d,  $J = 7.2$  Hz, 3H), 2.08 (s, 3H), 2.42 (s, 3H), 6.24 (q,  $J = 7.2$  Hz, 1H), 7.09-7.42 (m, 5H), 8.48 (m, 2H), 8.70 (m, 3H); MS (ES): 345.1 (M<sup>+</sup>+1).

- dl*-5,6-dimethyl-2-(2-furyl)-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ  
25 1.98 (d,  $J = 7.8$  Hz, 3H), 1.99 (s, 3H), 2.37 (s, 3H), 6.12 (q,  $J = 7.8$  Hz, 1H), 6.48 (dd,  $J = 1.8$  Hz, 3.6 Hz, 1H), 7.17-7.55 (m, 7H), 9.6 (s, 1H); MS (ES): 334.2 (M<sup>+</sup>+1).

- 30 *dl*-5,6-dimethyl-2-(3-furyl)-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.99 (d,  $J = 7$  Hz, 3H), 2.02 (s, 3H), 2.42 (s, 3H), 6.24 (q,  $J = 7$  Hz, 1H), 7.09 (s, 1H), 7.18-7.32 (m, 5H), 7.48 (s, 1H), 8.51 (s, 1H); MS (ES): 334.2 (M<sup>+</sup>+1).

35

*dl*-5,6-dimethyl-2-cyclopentyl-7*H*-7-(1-phenylethyl)

pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.95 (d,  $J = 7.4$  Hz, 3H), 2.00 (s, 3H), 2.33 (s, 3H), 1.68-1.88 (m, 8H), 2.97 (m, 1H), 6.10 (q,  $J = 7.4$  Hz, 1H), 7.16-7.30 (m, 5H), 9.29 (s, 1H); MS (ES): 336.3 ( $\text{M}^+ + 1$ ).

5

dl-5,6-dimethyl-2-(2-thienyl)-7H-7-(1-phenylethyl) pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.02 (d,  $J = 7.2$  Hz, 3H), 2.06 (s, 3H), 2.41 (s, 3H), 6.13 (q,  $J = 7.2$  Hz, 1H), 7.12 (dd,  $J = 4.8, 2.8$  Hz, 1H), 7.26-7.32 (m, 5H), 7.44 (d,  $J = 4.8$  Hz, 1H), 8.01 (d,  $J = 2.8$  Hz, 1H) 11.25 (s, 1H); MS (ES): 350.2 ( $\text{M}^+ + 1$ ).

dl-5,6-dimethyl-2-(3-thienyl)-7H-7-(1-phenylethyl) pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.00 (d,  $J = 7.4$  Hz, 3H), 2.05 (s, 3H), 2.43 (s, 3H), 6.24 (q,  $J = 7.4$  Hz, 1H), 7.24-7.33 (m, 5H), 7.33-7.39 (m, 1H), 7.85 (m, 1H), 8.47 (m, 1H), 12.01 (s, 1H); MS (ES): 350.2 ( $\text{M}^+ + 1$ ).

dl-5,6-dimethyl-2-(4-fluorophenyl)-7H-7-(1-phenylethyl) pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.01 (d,  $J = 6.8$  Hz, 3H), 2.05 (s, 3H), 2.42 (s, 3H), 6.26 (q,  $J = 6.8$  Hz, 1H), 7.12-7.36 (m, 7H), 8.23-8.30 (m, 2H), 11.82 (s, 1H); MS (ES): 362.3 ( $\text{M}^+ + 1$ ).

dl-5,6-dimethyl-2-(3-fluorophenyl)-7H-7-(1-phenylethyl) pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.02 (d,  $J = 7.4$  Hz, 3H), 2.06 (s, 3H), 2.44 (s, 3H), 6.29 (q,  $J = 7.4$  Hz, 1H), 7.13-7.51 (m, 7H), 8.00-8.04 (m, 2H), 11.72 (s, 1H); MS (ES): 362.2 ( $\text{M}^+ + 1$ ).

30

dl-5,6-dimethyl-2-(2-fluorophenyl)-7H-7-(1-phenylethyl) pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.00 (d,  $J = 7.2$  Hz, 3H), 2.05 (s, 3H), 2.38 (s, 3H), 6.24 (q,  $J = 7.2$  Hz, 1H), 7.18 - 7.45 (m, 8 H), 8.21 (m, 1H), 9.54 (s, 1H); MS (ES): 362.2 ( $\text{M}^+ + 1$ ).

35

*dl*-5,6-dimethyl-2-isopropyl-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one.

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.30 (d, *J* = 6.8 Hz, 3H), 1.32 (d, *J* = 7.0 Hz, 3H), 2.01 (s, 3H), 2.34 (s, 3H), 2.90 (m, 1H),  
5 6.13 (m, 1H), 7.17-7.34 (m, 5H), 10.16 (s, 1H), MS (ES): 310.2 (M<sup>+</sup>+1).

#### Preparation 8:

A solution of *dl*-1-(1-phenylethyl)-2-benzoylamino-3-cyano-4-  
10 dimethylpyrrole (785 mg, 2.38 mmol) with concentrated H<sub>2</sub>SO<sub>4</sub> (1 mL) in DMF (13 mL) was stirred at 130°C for 48 h. The black solution was diluted with CHCl<sub>3</sub> (100 mL) and washed with 1 N NaOH (30 mL), and brine (30 mL). The organic fraction was dried, filtered, concentrated, and purified by flash  
15 chromatography (SiO<sub>2</sub>; 8/2 EtOAc/Hex, *R<sub>f</sub>* 0.35) to a brown solid (184 mg, 24%) as *dl*-5-methyl-2-phenyl-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 8.18 (m, 2H, Ar-H), 7.62-7.44 (m, 3H, Ar-H), 7.40-7.18 (m, 5H, Ar-H), 6.48 (s, 1H, pyrrole-H), 6.28 (q, 1H, *J*  
20 = 7.2 Hz, CH-CH<sub>3</sub>), 2.18 (s, 3H, pyrrole-CH<sub>3</sub>), 2.07 (d, 3H, *J* = 7.2 Hz, CH-CH<sub>3</sub>); MS (ES): 330.2 (M<sup>+</sup> + 1).

#### Preparation 9:

A mixture of *dl*-1-(1-phenylethyl)-2-amino-3-cyano-4,5-  
25 dimethylpyrrole (9.60 g, 40.0 mmol) and of formic acid (50.0 mL, 98%) was refluxed for 5 hr. After cooling down to room temperature and scratching the sides of flask, copious precipitate was formed and filtered. The material was washed with water until washings showed neutral pH to give *dl*-5,6-  
30 dimethyl-7*H*-7-(1-phenylethyl)pyrrolo[2,3*d*]pyrimidin-4(3*H*)-one. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.96 (d, *J* = 7.4 hz, 3H), 2.00 (s, 3H), 2.38 (s, 3H), 6.21 (q, *J* = 7.4 Hz, 1H), 7.11-7.35 (m, 5H), 7.81 (s, 1H), 11.71 (s, 1H); MS (ES): 268.2 (M<sup>+</sup>+1).

#### 35 Preparation 10:

dl-5,6-dimethyl-2-phenyl-7H-7-(1-phenylethyl) pyrrolo  
[2,3d]pyrimidin-4(3H)-one (1.0 g, 2.91 mmol) was suspended in  
polyphosphoric acid (30.0 mL). The mixture was heated at 100°C  
for 4 hr. The hot suspension was poured onto ice water,  
5 stirred vigorously to disperse suspension, and basified to pH  
6 with solid KOH. The resulting solid was filtered and  
collected to give 0.49 g (69%) of 5,6-dimethyl-2-phenyl-7H-  
pyrrolo[2,3d]pyrimidin-4(3H)-one. <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>)  
δ 2.17 (s, 3H), 2.22 (s, 3H), 7.45 (br, 3H), 8.07 (br, 2H),  
10 11.49 (s, 1H), 11.82 (s, 1H); MS (ES): 344.2 (M<sup>+</sup>+1).

The following compounds were obtained in a similar manner as  
that of Preparation 10:

15 5-methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one. MS  
(ES): 226.0 (M<sup>+</sup>+1).

5,6-dimethyl-2-(3-pyridyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-  
one. MS (ES): 241.1 (M<sup>+</sup>+1).

20 5,6-dimethyl-2-(2-furyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  
<sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 2.13 (s, 3H), 2.18 (s, 3H), 6.39  
(dd, J = 1.8, 3.6 Hz, 1H), 6.65 (dd, J = 1.8 Hz, 3.6 Hz, 1H),  
7.85 (dd, J = 1.8, 3.6 Hz, 1H), 11.45 (s, 1H), 11.60 (s,  
25 1H); MS (ES): 230.1 (M<sup>+</sup>+1).

5,6-dimethyl-2-(3-furyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  
<sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 2.14 (s, 3H), 2.19 (s, 3H), 6.66  
(s, 1H), 7.78 (s, 1H), 8.35 (s, 1H), 11.3 (s, 1H), 11.4 (s,  
30 1H); MS (ES): 230.1 (M<sup>+</sup>+1).

5,6-dimethyl-2-cyclopentyl-7H-pyrrolo[2,3d]pyrimidin-4(3H)-  
one. <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 1.57-1.91 (m, 8 H), 2.12 (s,  
3H), 2.16 (s, 3H), 2.99 (m, 1H), 11.24 (s, 1H), 11.38 (s,  
35 1H); MS (ES): 232.2 (M<sup>+</sup>+1):

5,6-dimethyl-2-(2-thienyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.14 (s, 3H), 2.19 (s, 3H), 7.14 (dd,  $J$  = 3.0, 5.2 Hz, 1H), 7.70 (d,  $J$  = 5.2 Hz 1H), 8.10 (d,  $J$ =3.0 Hz, 1H), 11.50 (s, 1H); MS (ES): 246.1 ( $M^+$ +1).

5

5,6-dimethyl-2-(3-thienyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.17 (s, 3H), 2.21 (s, 3H), 7.66 (m, 1H), 7.75 (m, 1H), 8.43 (m, 1H), 11.47 (s, 1H), 11.69 (s, 1H); MS (ES): 246.1 ( $M^+$ +1).

10

5,6-dimethyl-2-(4-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.17 (s, 3H), 2.21 (s, 3H), 7.31 (m, 2H), 8.12 (m, 2H), 11.47 (s, 1H); MS (ES): 258.2 ( $M^+$ +1).

15

5,6-dimethyl-2-(3-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.18 (s, 3H), 2.21 (s, 3H), 7.33 (m, 1H), 7.52 (m, 1H), 7.85-7.95 (m, 2H), 11.56 (s, 1H), 11.80 (s, 1H); MS (ES): 258.1 ( $M^+$ +1).

20

5,6-dimethyl-2-(2-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.18 (s, 3H), 2.22 (s, 3H), 7.27-7.37 (m, 2H), 7.53 (m 1H), 7.68 (m, 1H), 11.54 (s, 1H), 11.78 (s, 1H); MS (ES): 258.1 ( $M^+$ +1).

25

5,6-dimethyl-2-isopropyl-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  1.17 (d,  $J$ = 6.6 Hz, 6H), 2.11 (s, 3H), 2.15 (s, 3H), 2.81 (m, 1H), 11.20 (s, 1H), 11.39 (s, 1H); MS (ES): 206.1 ( $M^+$ +1).

30

5,6-dimethyl-7H-pyrrolo[2,3d]pyrimidin-4(3H)-one.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.13 (s, 3H), 2.17 (s, 3H), 7.65 (s, 1H); MS (ES): 164.0 ( $M^+$ +1).

35 **Preparation 11:**

A solution of 5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]



pyrimidin-4(3H)-one (1.0 g, 4.2 mmol) in phosphorus  
oxychloride (25.0 mL) was refluxed for 6 hr and then  
concentrated in vacuo to dryness. Water was added to the  
residue to induce crystallization and the resulting solid was  
5 filtered and collected to give 0.90 g (83%) of 4-chloro-5,6-  
dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz,  
DMSO-d<sub>6</sub>) δ 2.33 (s, 3H), 2.33 (s, 3H), 7.46-7.49 (m, 3H),  
8.30-8.35 (m, 2H), 12.20 (s, 1H); MS (ES): 258.1 (M<sup>+</sup>+1).  
The following compounds were obtained in a similar manner as  
10 that of Preparation 11:

4-chloro-5-methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS  
(ES): 244.0 (M<sup>+</sup>+1).  
15 4-chloro-6-methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS  
(ES): 244.0 (M<sup>+</sup>+1).

4-chloro-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz,  
DMSO-d<sub>6</sub>) 8.35 (2, 2H), 7.63 (br s, 1H), 7.45 (m, 3H), 6.47  
20 (br s, 1H); MS (ES): 230.0 (M<sup>+</sup>+1).

4-chloro-5,6-dimethyl-2-(3-pyridyl)-7H-pyrrolo[2,3d]  
pyrimidine. MS (ES): 259.0 (M<sup>+</sup>+1).

25 4-chloro-5,6-dimethyl-2-(2-furyl)-7H-pyrrolo[2,3d]pyrimidine.  
<sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 2.35 (s, 3H), 2.35 (s, 3H), 6.68  
(dd, J = 1.8, 3.6 Hz, 1H), 7.34 (dd, J = 1.8 Hz, 3.6 Hz, 1H),  
7.89 (dd, J = 1.8, 3.6 Hz, 1H); MS (ES): 248.0 (M<sup>+</sup>+1).

30 4-chloro-5,6-dimethyl-2-(3-furyl)-7H-pyrrolo[2,3d]pyrimidine.  
<sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 2.31 (s, 3H), 2.31 (s, 3H), 6.62  
(s, 1H), 7.78 (s, 1H), 8.18 (s, 1H), 12.02 (s, 1H); MS (ES):  
248.1 (M<sup>+</sup>+1).

35 4-chloro-5,6-dimethyl-2-cyclopentyl-7H-pyrrolo[2,3d]  
pyrimidine. <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 1.61- 1.96 (m, 8H),

2.27 (s, 3H), 2.27 (s, 3H), 3.22 (m, 1H), 11.97 (s, 1H); MS (ES): 250.1 ( $M^+ + 1$ ).

- 4-chloro-5,6-dimethyl-2-(2-thienyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.29 (s, 3H), 2.31 (s, 3H), 7.14 (dd,  $J = 3.1$  Hz, 4.0 Hz, 1H), 7.33 (d,  $J = 4.9$  Hz, 1H), 7.82 (d,  $J = 3.1$  Hz, 1H), 12.19 (s, 1H); MS (ES): 264.1 ( $M^+ + 1$ ).
- 10 4-chloro-5,6-dimethyl-2-(3-thienyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.32 (s, 3H), 2.32 (s, 3H), 7.62 (dd,  $J = 3.0$ , 5.2 Hz, 1H), 7.75 (d,  $J = 5.2$  Hz, 1H), 8.20 (d,  $J = 3.0$  Hz, 1H); MS (ES): 264.0 ( $M^+ + 1$ ).
- 15 4-chloro-5,6-dimethyl-2-(4-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.33 (s, 3H), 2.33 (s, 3H), 7.30 (m, 2H), 8.34 (m, 2H), 12.11 (s, 1H); MS (ES): 276.1 ( $M^+ + 1$ ).
- 20 4-chloro-5,6-dimethyl-2-(3-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.31 (s, 3H), 2.33 (s, 3H), 7.29 (m, 1H), 7.52 (m, 1H), 7.96 (m, 1H), 8.14 (m, 1H), 11.57 (s, 1H); MS (ES): 276.1 ( $M^+ + 1$ ).
- 25 4-chloro-5,6-dimethyl-2-(2-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.34 (s, 3H), 2.34 (s, 3H), 7.33 (m, 2H), 7.44 (m, 1H), 7.99 (m, 1H), 12.23 (s, 1H); MS (ES): 276.1 ( $M^+ + 1$ ).
- 30 4-chloro-5,6-dimethyl-2-isopropyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  1.24 (d,  $J = 6.6$  Hz, 6H), 2.28 (s, 3H), 2.28 (s, 3H), 3.08 (q,  $J = 6.6$  Hz, 1H), 11.95 (s, 1H); MS (ES): 224.0 ( $M^+ + 1$ ).
- 35 4-chloro-5,6-dimethyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz, DMSO- $d_6$ )  $\delta$  2.31 (s, 3H), 2.32 (s, 3H), 8.40 (s, 1H); MS (ES): 182.0 ( $M^+ + 1$ ).

*dl*-4-chloro-5,6-dimethyl-2-phenyl-7*H*-7-(1-phenylethyl)pyrrolo  
[2,3*d*]pyrimidine.

**Preparation 12:**

- 5 To a solution of *dl*-1,2-diaminopropane (1.48 g, 20.0 mmol)  
and sodium carbonate (2.73 g, 22.0 mmol) in dioxane (100.0  
mL) and water (100.0 mL) was added di-*tert*-dicarbonate (4.80  
g, 22.0 mmol) at room temperature. The resulted mixture was  
stirred for 14 hr. Dioxane was removed in *vacuo*. The  
10 precipitate was filtered off and the filtrate was  
concentrated in *vacuo* to dryness. The residue was triturated  
with EtOAc and then filtered. The filtrate was concentrated  
in *vacuo* to dryness to give a mixture of *dl*-1-amino-2-(1,1-  
dimethylethoxy)carbonylamino-propane and *dl*-2-amino-1-(1,1-  
15 dimethylethoxy)carbonylamino-propane which were not separable  
by normal chromatography method. The mixture was used for the  
reaction in Example 8.

**Preparation 13:**

- 20 To solution of Fmoc- $\beta$ -Ala-OH (1.0 g, 3.212 mmol) and oxalyl  
chloride (0.428 g, 0.29 mL, 3.373 mmol) in dichloromethane  
(20.0 mL) was added a few drops of *N,N*-dimethylformamide at  
0°C. The mixture was stirred at room temperature for 1 hr  
followed by addition of cyclopropylmethylamine (0.229 g, 0.28  
25 mL, 3.212 mmol) and triethylamine (0.65 g, 0.90 mL, 6.424  
mmol). After 10 min, the mixture was treated with 1 M  
hydrochloride (10.0 mL) and the aqueous mixture was extracted  
with dichloromethane (3 x 30.0 mL). The organic solution was  
concentrated in *vacuo* to dryness. The residue was treated  
30 with a solution of 20% piperidine in *N,N*-dimethylformamide  
(20.0 mL) for 0.5 hr. After removal of the solvent in *vacuo*,  
the residue was treated with 1 M hydrochloride (20.0 mL) and  
ethyl acetate (20.0 mL). The mixture was separated and the  
aqueous layer was basified with solid sodium hydroxide to pH  
35 = 8. The precipitate was removed by filtration and the  
aqueous solution was subjected to ion exchange column eluted

with 20% pyridine to give 0.262 g (57%) of N-cyclopropylmethyl  $\beta$ -alanine amide.  $^1\text{H}$  NMR (200 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  0.22 (m, 2H), 0.49 (m, 2H), 0.96 (m, 2H), 2.40 (t, 2H), 2.92 (t, 2H), 3.05 (d, 2H); MS (ES): 143.1 ( $\text{M}^+ + 1$ ).

5

#### Preparation 14:

N-tert-butoxycarbonyl-trans-1,4-cyclohexyldiamine.

trans-1,4-cyclohexyldiamine (6.08 g, 53.2 mmol) was dissolved in dichloromethane (100mL). A solution of di-t-butylidicarbonate (2.32 g, 10.65 mmol in 40 mL dichloromethane) was added via cannula. After 20 hours, the reaction was partitioned between  $\text{CHCl}_3$  and water. The layers were separated and the aqueous layer was extracted with  $\text{CHCl}_3$  (3x). The combined organic layers were dried over  $\text{MgSO}_4$ ,  
10 filtered and concentrated to yield 1.20 g of a white solid (53%).  $^1\text{H}$ -NMR (200MHz,  $\text{CDCl}_3$ ):  $\delta$  1.0-1.3 (m, 4H), 1.44 (s, 9H), 1.8-2.1 (m, 4H), 2.62 (brm, 1H), 3.40 (brs, 1H), 4.37 (brs, 1H); MS (ES): 215.2 ( $\text{M}^+ + 1$ ).

20 4-(N-acetyl)-N-tert-butoxycarbonyl-trans-1,4-cyclohexyldiamine.

N-tert-butoxycarbonyl-trans-1,4-cyclohexyldiamine (530 mg, 2.47 mmol) was dissolved in dichloromethane (20 mL). Acetic anhydride (250 mg, 2.60 mmol) was added dropwise. After 16  
25 hours, the reaction was diluted with water and  $\text{CHCl}_3$ . The layers were separated and the aqueous layer was extracted with  $\text{CHCl}_3$  (3x). The combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated. Recrystallization (EtOH/ $\text{H}_2\text{O}$ ) yielded 190 mg of white crystals (30%).  $^1\text{H}$  NMR  
30 (200 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.9 - 1.30 (m, 4H), 1.43 (s, 9H), 1.96-2.10 (m, 7H), 3.40 (brs, 1H), 3.70 (brs, 1H), 4.40 (brs, 1H), 4.40 (brs, 1H); MS (ES): 257.2 ( $\text{M}^+ + 1$ ), 242.1 ( $\text{M}^+ - 15$ ), 201.1 ( $\text{M}^+ - 56$ ).

35 4-(4-trans-acetamidocyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-(1-phenylethyl) pyrrolo[2,3d]pyrimidine.

4-(N-acetyl)-N-tert-butoxycarbonyl-trans-1,4-

cyclohexyldiamine (190 mg, 0.74 mmol), was dissolved in dichloromethane (5 mL) and diluted with TFA (6 ml). After 16 hours, the reaction was concentrated. The crude solid, DMSO (2mL), NaHCO<sub>3</sub> (200 mg, 2.2 mmol) and 4-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (35 mg, 0.14 mmol) were combined in a flask and heated to 130 °C. After 4.5 hours, the reaction was cooled to room temperature and diluted with EtOAc and water. The layers were separated and the aqueous layer was extracted with EtOAc (3x). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated. Chromatography (silica preparatory plate; 20:1 CHCl<sub>3</sub>:EtOH) yielded 0.3 mg of a tan solid (1% yield). MS (ES): 378.2 (M<sup>+</sup>+1).

4 - (N-methanesulfonyl)-N-tert-butoxycarbonyl-trans-1,4-cyclohexyldiamine.  
trans-1,4-cyclohexyldiamine (530 mg, 2.47 mmol) was dissolved in dichloromethane (20 ml) and diluted with pyridine (233 mg, 3.0 mmol). Methanesulfonyl chloride (300 mg, 2.60 mmol) was added dropwise. After 16 hours, the reaction was diluted with water and CHCl<sub>3</sub>. The layers were separated and the aqueous layer was extracted with CHCl<sub>3</sub> (3x). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated. recrystallization (EtOH/H<sub>2</sub>O) yielded 206 mg of white crystals (29%). <sup>1</sup>H-NMR (200MHz, CDCl<sub>3</sub>): δ 1.10-1.40 (m, 4H), 1.45 (s, 9H), 2.00-2.20 (m, 4H), 2.98 (s, 3H), 3.20-3.50 (brs, 2H), 4.37 (brs, 1H); MS (ES) 293.1 (M<sup>+</sup>+1), 278.1 (M<sup>+</sup>-15), 237.1 (M<sup>+</sup>-56).

4 - (4-trans-methanesulfamidocyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-(1-phenylethyl)pyrrolo[2,3d]pyrimidine.  
4 - (N-sulfonyl)-N-tert-butoxycarbonyl-trans-1,4-cyclohexyldiamine (206 mg, 0.71 mmol), was dissolved in dichloromethane (5ml) and diluted with TFA (6 ml). After 16 hours, the reaction was concentrated. The crude reaction mixture, DMSO (2 ml), NaHCO<sub>3</sub> (100 mg, 1.1 mmol) and 1-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine were

combined in a flask and heated to 130 °C. After 15 hours, the reaction was cooled to room temperature, and diluted with EtOAc (3x). The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated. Chromatography (silica preparatory plate, 20:1 CHCl<sub>3</sub>/EtOH) yielded 2.6 mg of a tan solid (5% yield). MS (ES): 414.2 (M<sup>+</sup>+1).

**Example 1:**

A solution of 4-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (0.50 g, 1.94 mmol) and 4-trans-hydroxycyclohexylamine (2.23 g, 19.4 mmol) in methyl sulfoxide (10.0 mL) was heated at 130°C for 5 hr. After cooling down to room temperature, water (10.0 mL) was added and the resulted aqueous solution was extracted with EtOAc (3 x10.0 mL). The combined EtOAc solution was dried (MgSO<sub>4</sub>) and filtered, the filtrate was concentrated in vacuo to dryness, the residue was chromatographed on silica gel to give 0.49 g (75%) of 4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. mp 197-199°C; <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ<sub>1.25-1.59</sub> (m, 8H), 2.08 (s, 3H), 2.29 (s, 3H), 3.68-3.79 (m, 1H), 4.32-4.38 (m, 1H), 4.88 (d, J = 8 Hz, 1H), 7.26-7.49 (m, 3H), 8.40-8.44 (dd, J = 2.2, 8 Hz, 2H), 10.60 (s, 1H); MS (ES): 337.2 (M<sup>+</sup>+1).

The following compounds were obtained in a similar manner to that of Example 1:

4-(4-trans-hydroxycyclohexyl)amino-6-methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ<sub>11.37</sub> (s, 1H, pyrrole-NH), 8.45 (m, 2H, Ar-H), 7.55 (m, 3H, Ar-H), 6.17 (s, 1H, pyrrole-H), 4.90 (br d, 1H, NH), 4.18 (m, 1H, CH-O), 3.69 (m, 1H, CH-N), 2.40-2.20 (m, 2H), 2.19-1.98 (m, 2H), 2.25 (s, 3H, CH<sub>3</sub>) 1.68-1.20 (m, 4H); MS (ES): 323.2 (M<sup>+</sup>+1).

4-(4-trans-hydroxycyclohexyl)amino-5-methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ<sub>11.37</sub> (s, 1H, pyrrole-NH), 8.40 (m, 2H, Ar-H), 7.45 (m, 3H, Ar-H), 5.96

(s, 1H, pyrrole-H), 4.90 (br d, 1H, NH), 4.18 (m, 1H, CH-O), 3.69 (m, 1H, CH-N), 2.38-2.20 (m, 2H), 2.18-1.98 (m, 2H), 2.00 (s, 3H, CH<sub>3</sub>) 1.68-1.20 (m, 4H); MS (ES): 323.2 (M<sup>+</sup>+1).

5 4-(4-trans-hydroxycyclohexyl)amino-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. mp 245.5-246.5°C; <sup>1</sup>H NMR (200MHz, CD<sub>3</sub>OD) δ 8.33 (m, 2H, Ar-H), 7.42 (m, 3H, Ar-H), 7.02 (d, 1H, J=3.6 Hz, pyrrole-H), 6.53 (d, 1H, J=3.6 Hz, pyrrole-H), 4.26 (m, 1H, CH-O), 3.62 (m, 1H, CH-N), 2.30-2.12 (m, 2H), 2.12-1.96 (m, 10 2H), 1.64-1.34 (m, 4H); MS, M+1=309.3; Anal (C<sub>19</sub>H<sub>22</sub>N<sub>4</sub>O) C, H, N.

4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-(3-pyridyl)-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.21-1.54 (m, 8H); 2.28 (s, 3H); 2.33 (s, 3H); 3.70 (m, 1H), 4.31 (m, 1H), 4.89 (d, 1H), 7.40 (m, 1H), 8.61 (m, 2H), 9.64 (m, 1H); MS (ES): 338.2 (M<sup>+</sup>+1).

4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-(2-furyl)-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.26-1.64 (m, 8H), 2.22 (s, 3H), 2.30 (s, 3H), 3.72 (m, 1H), 4.23 (m, 1H), 4.85 (d, 1H), 6.52 (m, 1H), 7.12 (m, 1H), 7.53 (m, 1H), 9.28 (s, 1H); MS (ES): 327.2 (M<sup>+</sup>+1).

25 4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-(3-furyl)-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.25-1.63 (m, 8 H), 2.11 (s, 3H), 2.27 (s, 3H), 3.71 (m, 1H), 4.20 (m, 1H), 4.84 (d, 1H), 7.03 (m, 1H), 7.45 (m, 1H), 8.13 (m, 1H), 10.38 (m, 1H); MS (ES): 327.2 (M<sup>+</sup>+1).

30 4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-cyclopentyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.26-2.04 (m, 16 H), 2.26 (s, 3H), 2.27 (s, 3H), 3.15 (m, 1H), 3.70 (m, 1H), 4.12 (m, 1H), 4.75 (d, 1H); MS 35 (ES): 329.2 (M<sup>+</sup>+1).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-(2-thienyl)-7H-pyrrolo[2,3d]pyrimidin-4-amine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.28-1.59 (m, 8H), 2.19 (s, 3H), 2.29 (s, 3H), 3.74 (m, 1H), 4.19 (m, 1H), 4.84 (d, 1H), 7.09 (m, 1H), 7.34 (m, 1H), 7.85 (m, 1H), 9.02 (s, 1H); MS (ES): 343.2 ( $\text{M}^+ + 1$ ).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-(3-thienyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.21-1.60 (m, 8H), 1.98 (s, 3H), 2.23 (s, 3H), 3.66 (m, 1H), 4.22 (m, 1H), 7.27 (m, 1H), 7.86 (m, 1H), 8.09 (m, 1H), 11.23 (s, 1H); MS (ES): 343.2 ( $\text{M}^+ + 1$ ).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-(4-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.26-1.66 (m, 8H), 1.94 (s, 3H), 2.28 (s, 3H), 3.73 (m, 1H), 4.33 (m, 1H), 4.92 (d, 1H), 7.13 (m, 2H), 8.41 (m, 2H), 11.14 (s, 1H); MS (ES): 355.2 ( $\text{M}^+ + 1$ ).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-(3-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.26-1.71 (m, 8H), 2.06 (s, 3H), 2.30 (s, 3H), 3.72 (m, 1H), 4.30 (m, 1H), 4.90 (d, 1H), 7.09 (m, 1H), 7.39 (m, 1H), 8.05 (m, 1H), 8.20 (m, 1H), 10.04 (s, 1H); MS (ES): 355.2 ( $\text{M}^+ + 1$ ).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-(2-fluorophenyl)-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.30-1.64 (m, 8H), 2.17 (s, 3H), 2.31 (s, 3H), 3.73 (m, 1H), 4.24 (m, 1H), 4.82 (d, 1H), 7.28 (m, 2H), 8.18 (m, 1H), 9.02 (m, 1H), 12.20 (s, 1H); MS (ES): 355.3 ( $\text{M}^+ + 1$ ).

4-(4-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-isopropyl-7H-pyrrolo[2,3d]pyrimidine  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.31 (d,  $J = 7.0$  Hz, 6H), 1.30-1.65 (m, 8H), 2.27 (s, 3H), 2.28 (s, 3H), 3.01 (m,  $J = 7.0$  Hz, 1H), 3.71 (m, 1H), 4.14 (m, 1H), 4.78 (d, 1H); MS (ES): 303.2.



dl-4-(2-*trans*-hydroxycyclohexyl)amino-5,6-dimethyl-2-isopropyl-7H-pyrrolo[2,3d]pyrimidine <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) d 1.31-1.42 (br, 4H), 1.75-1.82 (br, 4H), 2.02 (s, 3H), 2.29 (s, 3H), 3.53 (m, 1H), 4.02 (m, 1H), 5.08 (d, 1H), 7.41-7.48 (m, 3H), 8.30 (m, 2H), 10.08 (s, 1H); MS (ES): 337.2 (M<sup>+</sup>+1).

4-(3,4-*trans*-dihydroxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 353.2 (M<sup>+</sup>+1).

10 4-(3,4-*cis*-dihydroxylcyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 353.2 (M<sup>+</sup>+1).

4-(2-acetyl aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.

15 mp 196-199°C; <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.72 (s, 3H), 1.97 (s, 3H), 2.31 (s, 3H), 3.59 (m, 2H), 3.96 (m, 2H), 5.63 (br, 1H), 7.44-7.47 (m, 3H), 8.36-8.43 (dd, J = 1 Hz, 7 Hz, 2H), 10.76 (s, 1H); MS (ES): 324.5 (M<sup>+</sup>+1).

20 dl-4-(2-*trans*-hydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.<sup>1</sup>

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.62 (m, 2H), 1.79 (br, 4H), 1.92 (s, 3H), 2.29 (s, 3H), 4.11 (m, 1H), 4.23 (m, 1H), 5.28 (d, 1H), 7.41-7.49 (m, 3H), 8.22 (m, 2H), 10.51 (s, 1H); MS (ES): 323.2 (M<sup>+</sup>+1).

<sup>1</sup> For preparation of 2-*trans*-hydroxycyclopentylamine, see PCT 9417090.

30 dl-4-(3-*trans*-hydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.<sup>1</sup>

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.58-1.90 (br, 6 H), 2.05 (s, 3H), 2.29 (s, 3H), 4.48-4.57 (m, 1H), 4.91-5.01 (m, 2H), 7.35-7.46 (m, 3H), 8.42-8.47 (m, 2H), 10.11 (s, 1H); MS (ES): 323.2 (M<sup>+</sup>+1).

007645-49100  
007645-49100

For preparation of 3-trans-hydroxycyclopentylamine, see EP-A-322242.

5 dl-4-(3-cis-hydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.<sup>2</sup>

<sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.82-2.28 (br, 6H), 2.02 (s, 3H), 2.30 (s, 3H), 4.53-4.60 (m, 1H), 4.95-5.08 (m, 1H), 5.85-5.93 (d, 1H), 7.35-7.47 (m, 3H), 8.42-8.46 (m, 2H), 10.05 (s, 1H); MS (ES): 323.2 (M<sup>+</sup>+1).

10 For preparation of 3-cis-hydroxycyclopentylamine, see EP-A-322242.

4-(3,4-trans-dihydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.<sup>1</sup> <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.92-1.99 (br, 2H), 2.14 (s, 3H), 2.20 (br, 2H), 2.30 (s, 3H), 2.41-2.52 (br, 2H), 4.35 (m, 2H), 4.98 (m, 2H), 7.38-7.47 (m, 3H), 8.38-8.42 (m, 2H), 9.53 (s, 1H); MS (ES): 339.2 (M<sup>+</sup>+1).

For preparation of 3,4-trans-dihydroxycyclopentylamine, see PCT 9417090.

20

4-(3-amino-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.

25 <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 2.02 (s, 3H), 2.29 (s, 3H), 2.71 (t, 2H), 4.18 (m, 2H), 5.75-5.95 (m, 3H), 7.38-7.48 (m, 3H), 8.37-8.41 (m, 2H), 10.42 (s, 1H); MS (ES): 310.1 (M<sup>+</sup>+1).

4-(3-N-cyclopropylmethylamino-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 0.51 (q, 2H), 0.40 (q, 2H), 1.79-1.95 (br, 1H), 2.36 (s, 3H), 2.40 (s, 3H), 2.72 (t, 2H), 2.99 (d, 2H), 4.04 (t, 2H), 7.58-7.62 (m, 3H), 8.22-8.29 (m, 2H); MS (ES): 364.2 (M<sup>+</sup>+1).

30

4-(2-amino-2-oxoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 2.31 (s, 3H), 2.38 (s, 3H), 4.26 (s, 2H), 7.36 (m, 3H), 8.33 (m, 2H); MS (ES): 396.1 (M<sup>+</sup>+1).

35

4-(2-N-methylamino-2-oxoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.99 (s, 3H), 2.17 (s, 3H), 2.82 (d, 3H), 4.39 (d, 2H), 5.76 (t, 1H), 6.71 (br, 1H), 7.41-7.48 (m, 3H), 8.40 (m, 2H), 10.66 (s, 1H); MS (ES): 310.1 (M<sup>+</sup>+1).

4-(3-tert-butyloxyl-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.45 (s, 9H), 1.96 (s, 3H), 2.29 (s, 3H), 2.71 (t, 2H), 4.01 (q, 2H), 5.78 (t, 1H), 7.41-7.48 (m, 3H), 8.22-8.29 (m, 2H); MS (ES): 367.2 (M<sup>+</sup>+1).

4-(2-hydroxyethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.92 (s, 3H), 2.29 (s, 3H), 3.81-3.98 (br, 4H), 5.59 (t, 1H), 7.39-7.48 (m, 3H), 8.37 (m, 2H), 10.72 (s, 1H); MS (ES): 283.1 (M<sup>+</sup>+1).

4-(3-hydroxypropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.84 (m, 2H), 1.99 (s, 3H), 2.32 (s, 3H), 3.62 (t, 2H), 3.96 (m, 2H), 3.35 (t, 1H), 7.39-7.48 (m, 3H), 8.36 (m, 2H), 10.27 (s, 1H); MS (ES): 297.2 (M<sup>+</sup>+1).

4-(4-hydroxybutyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.71-1.82 (m, 4H), 1.99 (s, 3H), 2.31 (s, 3H), 3.68-3.80 (m, 4H), 5.20 (t, 1H), 7.41-7.49 (m, 3H), 8.41 (m, 2H), 10.37 (s, 1H); MS (ES): 311.2 (M<sup>+</sup>+1).

4-(4-trans-acetylaminocyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.

4-(4-trans-methylsulfonylaminocyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.

4-(2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-7-(1-phenylethyl)pyrrolo[2,3d]pyrimidine.

4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-1-phenylethyl)pyrrolo[2,3d]pyrimidine.

4-(3-pyridylmethyl)amino-5,6-dimethyl-2-phenyl-7H-7-(1-phenylethyl)pyrrolo[2,3d]pyrimidine.

10 4-(2-methylpropyl)amino-5,6-dimethyl-2-phenyl-7H-7-(1-phenylethyl)pyrrolo[2,3d]pyrimidine.

#### Example 2:

To a stirred suspension of triphenylphosphine (0.047 g, 0.179 mmol) and benzoic acid (0.022 g, 0.179 mmol) in THF (1.0 mL) cooled to 0°C was added 4-(4-trans-hydroxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (0.05 g, 0.149 mmol) at 0°C. Diethyl azodicarboxylate (0.028 ml, 0.179 mmol) was then added dropwise over 10 minutes. The reaction was then allowed to warm to room temperature. After reaction was complete by TLC the reaction mixture was quenched with aqueous sodium bicarbonate (3.0 mL). The aqueous phase was separated and extracted with ether (2 X 5.0 mL). The organic extracts were combined, dried, and concentrated in vacuo to dryness. To the residue was added ether (2.0 mL) and hexane (5.0 mL) whereupon the bulk of the triphenylphosphine oxide was filtered off. Concentration of the filtrate gave a viscous oil which was purified by column chromatography (hexane:ethyl acetate=4:1) to give 5.0 mg (7.6%) of 4-(4-cis-benzoyloxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 441.3 ( $M^+ + 1$ ). The reaction also produced 50.0 mg (84%) of 4-(3-cyclohexenyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 319.2 ( $M^+ + 1$ ).

**Example 3:**

To a solution of 4-(4-*cis*-benzoyloxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (5.0 mg, 0.0114 mmol) in ethanol (1.0 mL) was added 10 drops of 2M sodium hydroxide. After 1 hr, the reaction mixture was extracted with ethyl acetate (3 x 5.0 mL) and the organic layer was dried, filtered and concentrated in vacuo to dryness. The residue was subjected to column chromatography (hexane:ethyl acetate=4:1) to give 3.6 mg (94%) of 4-(4-*cis*-hydroxycyclohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 337.2 ( $M^+ + 1$ ).

The following compounds were obtained in a similar manner as that of Example 3:

15

4-(3-N,N-dimethyl-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.01 (s, 3H), 2.31 (s, 3H), 2.73 (t, 2H), 2.97 (s, 6H), 4.08 (m, 2H), 6.09 (t, 1H), 7.41-7.48 (m, 3H), 8.43 (m, 2H), 10.46 (s, 1H); MS (ES): 338.2 ( $M^+ + 1$ ).

20

4-(2-formylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.26 (s, 3H), 2.37 (s, 3H), 3.59-3.78 (m, 2H), 3.88-4.01 (m, 2H), 5.48-5.60 (m, 1H), 7.38-7.57 (m, 3H), 8.09 (s, 1H), 8.30-8.45 (m, 2H), 8.82 (s, 1H); MS (ES): 310.1 ( $M^+ + 1$ ).

25

4-(3-acetylaminopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 338.2 ( $M^+ + 1$ ).

30

**Example 4:**

4-(3-tert-butyloxy-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (70.0 mg, 0.191 mmol) was dissolved in trifluoroacetic acid:dichloromethane (1:1, 5.0 mL). The resulting solution was stirred at room temperature for 1 hr. and then refluxed for 2 hr. After cooling down to room temperature, the mixture was concentrated in vacuo to dryness. The residue was subjected to preparative thin layer chromatography (EtOAc:hexane: AcOH=7:2.5:0.5) to give 40.0 mg (68%) of 4-(3-hydroxy-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CD<sub>3</sub>OD) δ 2.32 (s, 3H), 2.38 (s, 3H), 2.81 (t, 2H), 4.01 (t, 2H), 7.55 (m, 3H), 8.24 (m, 2H); MS (ES): 311.1 (M<sup>+</sup>+1).

15

The following compound was obtained in a similar manner as that of Example 4:

4-(3-aminopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 296.1 (M<sup>+</sup>+1), 279.1 (M<sup>+</sup>-NH<sub>3</sub>).

**Example 5:**

4-(3-hydroxy-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (50.0 mg, 0.161 mmol) was dissolved in a mixture of N,N-dimethylformamide (0.50 mL), dioxane (0.50 mL) and water (0.25 mL). To this solution was added methylamine (0.02 mL, 40% wt in water, 0.242 mmol), triethylamine (0.085 mL) and N,N,N',N'-tetramethyl uronium tetrafluoroborate (61.2 mg, 0.203 mmol). After stirring at room temperature for 10 min, the solution was concentrated and the residue was subjected to preparative thin layer chromatography (EtOAc) to give 35.0 mg (67%) of 4-(3-N-methyl-3-oxopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.92 (s, 3H), 2.30 (s, 3H), 2.65 (t, 2H), 4.08 (t, 2H), 5.90 (t, 1H), 6.12 (m, 1H), 7.45 (m, 3H), 8.41 (m, 2H), 10.68 (s, 1H); MS

35

(ES): 311.1 ( $M^+ + 1$ ).

The following compounds were obtained in a similar manner as that of Example 5:

5

4-(2-cyclopropanecarbonylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 350.2 ( $M^+ + 1$ ).

4-(2-isobutyrylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 352.2 ( $M^+ + 1$ ).

4-(3-propionylaminopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  1.00-1.08 (t, 3H), 1.71-2.03 (m, 4H), 2.08 (s, 3H), 2.37 (s, 3H), 3.26-3.40 (m, 2H), 3.79-3.96 (m, 2H), 5.53-5.62 (m, 1H), 6.17-6.33 (m, 1H), 7.33-7.57 (m, 3H), 8.31-8.39 (m, 2H), 9.69 (s, 1H); MS (ES): 352.2 ( $M^+ + 1$ ).

4-(2-methylsulfonylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  2.18 (s, 3H), 2.27 (s, 3H), 2.92 (s, 3H), 3.39-3.53 (m, 2H), 3.71-3.88 (m, 2H), 5.31-5.39 (m, 1H), 6.17-6.33 (m, 1H), 7.36-7.43 (m, 3H), 8.20-8.25 (m, 2H), 9.52 (s, 1H); MS (ES): 360.2 ( $M^+ + 1$ ).

25 **Example 6:**

A mixture of 4-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (0.70 g, 2.72 mmol) and 1,2-diaminoethane (10.0 mL, 150 mmol) was refluxed under inert atmosphere for 6 hr. The excess amine was removed in vacuo, the residue was washed sequentially with ether and hexane to give 0.75 g (98%) of 4-(2-aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES); 282.2 ( $M^+ + 1$ ), 265.1 ( $M^+ - NH_3$ ).

**Example 7:**

35 To a solution of 4-(2-aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (70.0 mg, 0.249 mmol) and triethylamine (50.4 mg, 0.498 mmol) in dichloromethane (2.0

mL) was added propionyl chloride (25.6 mg, 0.024 mL, 0.274 mmol) at 0°C. After 1 hr, the mixture was concentrated in vacuo and the residue was subjected to preparative thin layer chromatography (EtOAc) to give 22.0 mg (26%) of 4-(2-propionylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 338.2 (M<sup>+</sup>+1).

The following compounds were obtained in a similar manner as that of Example 7:

10

4-(2-N'-methylureaethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 2.13 (s, 3H), 2.32 (s, 3H), 3.53 (d, 3H), 3.55 (m, 2H), 3.88 (m, 2H), 4.29 (m, 1H), 5.68 (t, 1H), 5.84 (m, 1H), 7.42 (m, 3H), 8.36 (dd, 2H), 9.52 (s, 1H); MS (ES): 339.3 (M<sup>+</sup>+1).

15

4-(2-N'-ethylureaethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 353.2 (M<sup>+</sup>+1).

20 **Example 8:**

To a solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (41.1 mg, 0.215 mmol), dimethylaminopyridine (2.4 mg, 0.020 mmol) and pyruvic acid (18.9 mg, 0.015 mL, 0.215 mmol) in dichloromethane (2.0 mL) was added 4-(2-aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (55.0 mg, 0.196 mmol). The mixture was stirred at room temperature for 4 hr. Usual workup and column chromatography (EtOAc) then gave 10.0 mg (15%) of 4-(2'-pyruvylamidoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.. MS (ES): 352.2 (M<sup>+</sup>+1).

30

**Example 9:**

To a solution of 4-(2-aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (60.0 mg, 0.213 mmol) in dichloromethane (2.0 mL) was added N-trimethylsilyl isocyanate (43.3 mg, 0.051 mL, 0.320 mmol). The mixture was

35



stirred at room temperature for 3 hr followed by addition of aqueous sodium bicarbonate. After filtration through small amount of silica gel, the filtrate was concentrated in vacuo to dryness to give 9.8 mg (14%) of 4-(2-ureaethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 325.2 ( $M^+ + 1$ ).

The following compounds were obtained in a similar manner as that of Example 9:

10

dl-4-(2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.28-1.32 (d,  $J=8$  Hz, 3 H), 1.66 (s, 3H), 1.96 (s, 3H), 2.30 (s, 3H) 3.76-3.83 (m, 2H), 4.10-4.30 (m, 1H), 5.60-5.66 (t,  $J=6$  Hz, 1H), 7.40-7.51 (m, 3H), 8.36-8.43 (m, 2H), 10.83 (s, 1H); MS (ES): 338.2 ( $M^+ + 1$ ).

15

(R)-4-(2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.31 (d, 3H), 1.66 (s, 3H) 1.99 (s, 3H), 2.31 (s, 3H), 3.78-3.83 (m, 2H), 4.17-4.22 (m, 1H), 5.67 (t, 1H), 7.38-7.5 (m, 3H), 8.39 (m, 2H), 10.81 (s, 1H); MS (ES): 338.2 ( $M^+ + 1$ ).

20

(R)-4-(1-methyl-2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.41 (d, 3H), 1.68 (s, 3H), 2.21 (s, 3H), 2.34 (s, 3H), 3.46-3.52 (br, m, 2H), 4.73 (m, 1H), 5.22 (d, 1H), 7.41-7.46 (m, 3H), 8.36-8.40 (m, 2H), 8.93 (s, 1H); MS (ES): 338.2 ( $M^+ + 1$ ).

25

(S)-4-(2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  1.31 (d, 3H), 1.66 (s, 3H) 2.26 (s, 3H), 2.35 (s, 3H), 3.78-3.83 (m, 2H), 4.17-4.22 (m, 1H), 5.67 (t, 1H), 7.38-7.5 (m, 3H), 8.39 (m, 2H), 8.67 (s, 1H); MS (ES): 338.2 ( $M^+ + 1$ ).

30

(S)-4-(1-methyl-2-acetylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$

35

1.41 (d, 3H), 1.68 (s, 3H), 2.05 (s, 3H), 2.32 (s, 3H), 3.46-3.52 (m, 2H), 4.73 (m, 1H), 5.22 (d, 1H), 7.41-7.46 (m, 3H), 8.36-8.40 (m, 2H), 10.13 (s, 1H); MS (ES): 338.2 (M<sup>+</sup>+1).

5 **Example 10:**

Reaction of 4-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine with the mixture of dl-1-amino-2-(1,1-dimethylethoxy)carbonylamino-propane and dl-2-amino-1-(1,1-dimethylethoxy)carbonylamino-propane was run in a similar manner as that of Example 1. The reaction gave a mixture of dl-4-(1-methyl-2-(1,1-dimethylethoxy)carbonylamino)ethylamino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine and dl-4-(2-methyl-2-(1,1-dimethylethoxy)carbonylamino)ethylamino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine which were separated by column chromatography (EtOAc:hexanes=1:3). The first fraction was dl-4-(1-methyl-2-(1,1-dimethylethoxy)carbonylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine: <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.29 - 1.38 (m, 12 H), 1.95 (s, 3H), 2.31 (s, 3H) 3.34-3.43 (m, 2H), 4.62-4.70 (m, 1H), 5.36-5.40 (d, J=8 Hz, 1H), 5.53 (br, 1H), 7.37-7.49 (m, 3H), 8.37-8.44 (m, 2H), 10.75 (s, 1H). MS 396.3 (M<sup>+</sup>+1); The second fraction was dl-4-(2-(1,1-dimethylethoxy)carbonylaminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine: <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.26-1.40 (m, 12 H), 2.00 (s, 3H), 2.31 (s, 3H) 3.60-3.90 (m, 2H), 3.95-4.10 (m, 1H), 5.41-5.44 (d, J=6.0 Hz, 1H), 5.65 (br, 1H), 7.40-7.46 (m, 3H), 8.37-8.44 (m, 2H), 10.89 (s, 1H); MS (ES): 396.2 (M<sup>+</sup>+1).

30 The following compounds were obtained in a similar manner as that of Example 10:

(S,S)-4-(2-acetylaminoethyl)cyclohexylamino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.43 (m, 4 H), 1.60 (s, 3 H), 1.83 (m, 2 H), 2.18 (s, 3 H), 2.30 (m, 2 H), 2.32 (s, 3 H), 3.73 (br, 1H), 4.25 (br, 1H), 5.29 (d, 1H), 7.43-7.48 (m, 3H), 8.35-8.40 (m, 2H), 9.05 (s, 1 H).

4-(2-methyl-2-acetylaminopropyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.51 (s, 6H), 1.56 (s, 3H), 2.07 (s, 3H), 2.36 (s, 3H), 3.76 (d, 2H), 5.78 (t, 1H), 7.41-7.48 (m, 3H), 7.93 (s, 1H), 8.39 (m, 2H), 10.07 (s, 1H); MS (ES): 352.3 (M<sup>+</sup>+1).

**Example 11:**

dl-4-(1-methyl-2-(1,1-dimethylethoxy) carbonyl aminoethyl) amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (60.6 mg, 0.153 mmol) was treated with trifluoroacetic acid (0.5 mL) in dichloromethane (2.0 mL) for 14 hr. The organic solvent was removed in vacuo to dryness. The residue was dissolved in N,N-dimethylformamide (2.0 mL) and triethylamine (2.0 mL). To the solution at 0°C was added acetic anhydride (17.2 mg, 0.016, 0.169 mmol). The resulted mixture was stirred at room temperature for 48 hr and then concentrated in vacuo to dryness. The residue was subjected to preparative thin layer chromatography (EtOAc) to give 27.0 mg (52%) of dl-4-(1-methyl-2-acetyl aminoethyl) amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>) δ 1.38-1.42 (d, J=8 Hz, 3 H), 1.69 (s, 3H), 2.01 (s, 3H), 2.32 (s, 3H) 3.38-3.60 (m, 2H), 4.65-4.80 (m, 1H), 5.23-5.26 (d, J=6 Hz, 1H), 7.40-7.51 (m, 3H), 8.37-8.43 (m, 2H), 10.44 (s, 1H); MS (ES): 338.2 (M<sup>+</sup>+1).

**Example 12:**

(R,R)-4-(2-aminocyclohexyl) amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine, prepared in a similar manner as that of Example 1 from 4-chloro-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (0.15 g, 0.583 mmol) and (1R, 2R)-(-)-1,2-diaminocyclohexane (0.63 g, 5.517 mmol), was treated with triethylamine (0.726 g, 7.175 mmol) and acetic anhydride (0.325 g, 3.18 mmol) in N,N-dimethylformamide (10.0 mL) at room temperature for 2 hr. After removal of solvent in vacuo, ethyl acetate (10.0 mL) and water (10.0 mL) were added to the residue. The mixture was separated and the aqueous layer was

5

10

**Example 13:**

15  
20

## 25

30

35

oxohexyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 353.2 ( $M^+ + 1$ ).

**Example 15:**

- 5 A solution of 4-(2-aminoethyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine (70.0 mg, 0.249 mmol) and succinic anhydride (27.0 mg, 0.274 mmol) in dichloromethane (4.0 mL) with 1 drop of N,N-dimethylformamide was stirred at room temperature for 4 hr. The reaction mixture was extracted with  
10 20% sodium hydroxide (3 x 5.0 mL). The aqueous solution was acidified with 3 M hydrochloride to pH = 7.0. The whole mixture was extracted with ethyl acetate (3 x 10 mL). The combined organic solution was dried ( $MgSO_4$ ) and filtered. The filtrate was concentrated *in vacuo* to dryness to give 15.0 mg  
15 (16%) of 4-(7-hydroxy-3-aza-4,7-dioxoheptyl)amino-5,6-dimethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 382.2 ( $M^+ + 1$ ).

**Example 16:**

- 20 To 10 mL of dimethylformamide (DMF) at room temperature were added 700 mg of 4-*cis*-3-hydroxycyclopentyl)amino-2-phenyl-5,6-dimethyl-7H-pyrrolo[2,3d]pyrimidine followed by 455 mg of N-Boc glycine, 20 mg of N,N-dimethylaminopyridine (DMAP), 293 mg of hydroxybenzotriazole (HOBT) and 622 mg of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCl).  
25 The reaction mixture was left stirring overnight. DMF was then removed under reduced pressure and the reaction mixture was partitioned between 20mL of ethyl acetate and 50mL of water. The aqueous portion was extracted further with 2x20mL  
30 of ethyl acetate and the combined organic portions were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. Purification on silica gel, eluting with ethyl acetate/hexane gave 410 mg of the desired product: 4-(*cis*-3-(N-t-butoxycarbonyl-2-aminoacetoxy)  
35 cyclopentyl) amino-2-phenyl-5,6,-dimethyl-7H-pyrrolo[2,3d]pyrimidine, MS (ES) ( $M^+ + 1$ )=480.2. The ester was then treated

with 5 mL of 20% trifluoroacetic acid in dichloromethane at room temperature, left over night and then concentrated. Trituration with ethyl acetate gave 300 mg of an off white solid; 4-(*cis*-3-(2-aminoacetoxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine trifluoroacetic acid salt, MS (ES) ( $M^+ + 1$ ) = 380.1.

One skilled in the art will appreciate that the following compounds can be synthesized by the methods disclosed above:

10

4-(*cis*-3-hydroxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine MS (ES) ( $M^+ + 1$ ) = 323.1.

4-(*cis*-3-(2-aminoacetoxycyclopentyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidinetrifluoroacetic acid salt MS (ES) ( $M^+ + 1$ ) = 380.1.

4-(3-acetamido)piperidinyl-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine MS (ES) ( $M^+ + 1$ ) = 364.2.

4-(2-*N'*-methylureapropyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine, MS (ES) ( $M^+ + 1$ ) = 353.4.

4-(2-acetamidobutyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine, MS (ES) ( $M^+ + 1$ ) = 352.4.

4-(2-*N'*-methylureabutyl)amino-5,6-dimethyl-2-phenyl-7*H*-pyrrolo[2,3*d*]pyrimidine MS (ES) ( $M^+ + 1$ ) = 367.5

4-(2-aminocyclopropylacetamidoethyl)amino-2-phenyl-7*H*-pyrrolo[2,3*d*] pyrimidine MS (ES) ( $M^+ + 1$ ) = 309.1.

35

4-(*trans*-4-hydroxycyclohexyl)amino-2-(3-chlorophenyl)-7*H*-

pyrrolo[2,3d] pyrimidine MS (ES) ( $M^+ + 1$ ) = 342.8.

4-(trans-4-hydroxycyclohexyl)amino-2-(3-fluorophenyl)-7H-pyrrolo [2,3d] pyrimidine MS (ES) ( $M^+ + 1$ ) = 327.2.


5

4-(trans-4-hydroxycyclohexyl)amino-2-(4-pyridyl)-7H-pyrrolo[2,3d]pyrimidine MS (ES) ( $M^+ + 1$ ) = 310.2.

# Example 17

10

## Scheme IX




The pyrrole nitrogen of (7) (Scheme IX) was protected with di-*t*-butyldicarbonate under basic conditions to yield the corresponding carbamate (22). Radical bromination of (22) proceeded regioselectively to yield bromide (23). In general, compound (23) served as a key electrophilic intermediate for various nucleophilic coupling partners. Displacement of the alkyl bromide with sodium phenolate trihydrate yielded compound (24). Subsequent displacement of the aryl chloride and removal of the *t*-butyl carbamate

protecting group occurred in one step yielding desired compound (25).

Detailed Synthesis of Compounds (22)-(25) in Accordance with


5 Scheme IX



15

Di-t-butyl dicarbonate (5.37 g, 24.6 mmol) and dimethyl aminopyridine (1.13 g, 9.2 mmol) were added to a solution containing (7) (1.50 g, 6.15 mmol) and pyridine (30 mL). After 20 h the reaction was concentrated and the residue was  
20 partitioned between  $\text{CH}_2\text{Cl}_2$  and water. The  $\text{CH}_2\text{Cl}_2$  layer was separated, dried over  $\text{MgSO}_4$ , filtered and concentrated to yield a black solid. Flash chromatography ( $\text{SiO}_2$ ; 1/9 EtOAc/Hexanes,  $R_f$  0.40) yielded 1.70 g (80%) of a white solid (22).  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.50 (m, 2H, Ar-H), 7.45 (m, 3H, Ar-H), 6.39 (s, 1H, pyrrole-H), 2.66 (s, 3H, pyrrole- $\text{CH}_3$ ), 1.76 (s, 9H, carbamate- $\text{CH}_3$ ); MS,  $M + 1 = 344.1$ ; Mpt = 175-177°C.


30




35



N-Bromosuccinimide (508 mg, 2.86 mmol) and AIBN (112 mg, 0.68 mmol) were added to a solution containing (22) (935 mg, 2.71 mmol) and  $\text{CCl}_4$  (50 mL). The solution was heated to reflux. After 2 h the reaction was cooled to room temperature and concentrated *in vacuo* to yield a white solid. Flash chromatography ( $\text{SiO}_2$ ; 1/1  $\text{CH}_2\text{Cl}_2$ /Hexanes,  $R_f$  0.30) yielded 960 mg (84%) of a white solid (23).  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.52 (m, 2H, Ar-H), 7.48 (m, 3H, Ar-H), 6.76 (s, 1H, pyrrole-H), 4.93 (s, 2H, pyrrole- $\text{CH}_2\text{Br}$ ), 1.79 (s, 9H, carbamate- $\text{CH}_3$ ); MS,  $M + 1 = 423.9$ ; Mpt = 155-157°C.



Sodium phenoxide trihydrate (173 mg, 1.02 mmol) was added in one portion to a solution of bromide (23) (410 mg, 0.97 mmol) dissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (10 mL). After 2 h the reaction solution was partitioned between  $\text{CH}_2\text{Cl}_2$  and water. The water layer was extracted with  $\text{CH}_2\text{Cl}_2$ . The combined  $\text{CH}_2\text{Cl}_2$  layers were washed with water, dried over  $\text{MgSO}_4$ , filtered and concentrated to yield a yellow solid. Flash chromatography ( $\text{SiO}_2$ ; 1/6 EtOAc/Hexanes,  $R_f$  0.30) yielded 210 mg (50%) of a white solid (24).  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.53 (m, 2H, Ar-H), 7.48 (m, 3H, Ar-H), 7.34 (m, 2H, Ar-H), 7.03 (m, 3H, Ar-H), 6.83 (s, 1H, pyrrole-H), 5.45 (s, 2H, Ar $\text{CH}_2\text{O}$ ), 1.76 (s, 9H, carbamate- $\text{CH}_3$ ); MS,  $M^+ = 436.2$ .



A solution containing (24) (85 mg, 0.20 mmol), N-acetyleneethylenediamine (201 mg, 1.95 mmol) and DMSO (3 mL) was heated to 100°C. After 1 h the temperature was raised to 130°C. After 3 h the reaction was cooled to room temperature and partitioned between EtOAc and water. The water layer was extracted with EtOAc (2x). The combined EtOAc layers are washed with water, dried over MgSO<sub>4</sub>, filtered and concentrated. Flash chromatography (SiO<sub>2</sub>; 1/10 EtOH/ CHCl<sub>3</sub>, R<sub>f</sub> 0.25) yielded 73 mg (93%) of a white foamy solid (25). <sup>1</sup>H NMR (200 MHz, DMSO-d<sub>6</sub>) δ 11.81 (br s, 1H, N-H), 8.39 (m, 2H, Ar-H), 8.03 (br t, 1H, N-H), 7.57 (br t, 1H, N-H), 7.20 - 7.50 (m, 5H, Ar-H), 6.89 - 7.09 (m, 3H, Ar-H), 6.59 (s, 1H, pyrrole-H), 5.12 (s, 2H, ArCH<sub>2</sub>O), 3.61 (m, 2H, NCH<sub>2</sub>), 3.36 (m, 2H, NCH<sub>2</sub>), 1.79 (s, 3H, COCH<sub>3</sub>); MS, M+ 1 = 402.6

The following compounds were obtained in a manner similar to that of Example 17:

4-(2-acetylaminoethyl)amino-6-phenoxy-2-phenyl-7H-pyrrolo[2,3-d]pyrimidine. mp 196-197°C; MS (ES): 401.6 (M<sup>+</sup>+1).

4-(2-acetylaminoethyl)amino-6-(4-fluorophenoxy)methyl-2-phenyl-7H-pyrrolo[2,3-d]pyrimidine. MS(ES): 420.1 (M<sup>+</sup>+1).

4-(2-acetylaminoethyl)amino-6-(4-chlorophenoxy)methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS(ES): 436.1 ( $M^+ + 1$ ).

4-(2-acetylaminoethyl)amino-6-(4-methoxyphenoxy)methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS(ES): 432.1 ( $M^+ + 1$ ).

4-(2-acetylaminoethyl)amino-6-(N-pyridin-2-one)methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS(ES): 403.1 ( $M^+ + 1$ ).

10 4-(2-acetylaminoethyl)amino-6-(N-phenylamino)methyl-2-phenyl-7H-pyrrolo[2,3]pyrimidine. MS(ES): 400.9 ( $M^+ + 1$ ).

4-(2-acetylaminoethyl)amino-6-(N-methyl-N-phenylamino)methyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS(ES): 414.8 ( $M^+ + 1$ ).

15 4-(2-N'-methylureaethyl)amino-6-phenoxyethyl-2-phenyl-7H-pyrrolo[2,3d]pyrimidine. MS (ES): 416.9 ( $M^+ + 1$ ).

**Example 18: Adenosine A<sub>2</sub> Antagonist Experimentals**

20 Compound 1700 (Table 15 below): MS (ES): 366.1 ( $M^+ + 1$ ).

Compound 1710 (Table 15 below): MS (ES): 381.1 ( $M^+ + 1$ ).

Compound 1316 (Table 15 below): MS (ES): 353.2 ( $M^+ + 1$ )

25 Compound 1703 (Table 15 below): MS (ES): 357.1 ( $M^+ + 1$ ).

Compound 1719 (Table 15 below): <sup>1</sup>H-NMR (200MHz, d<sub>6</sub>-DMSO) (1.75 (m, 2H), 3.11 (m, 2H), 3.35 (s, 3H), 3.59 (m, 2H), 5.72 (m, 1H), 5.96 (m, 1H), 6.55 (s, 1H), 7.15 (s, 1H), 7.49 (m, 2H), 8.32 (m, 2H).

30

Compound 1704 (Table 15 below): MS (ES): 367.0 ( $M^+ + 1$ ).

Compound 1706 (Table 15 below): <sup>1</sup>H-NMR (200MHz, CDCl<sub>3</sub>) δ 1.22 (m, 2H), 1.60-2.40 (m, 4H), 4.53 (m, 1H), 4.94 (m, 1H), 5.70 (d, 1H, J = 8.2 Hz), 6.35 (d, 1H, J = 2.8 Hz), 6.97 (d, 1H, J = 2.0 Hz), 7.50 (m, 3H), 8.40 (m, 2H), 10.83 (brs, 1H).

35

Compound 1708 (Table 15 below): MS (ES) 399.0 (M<sup>+</sup>+1).

Compound 1709 (Table 15 below): MC (ES) 385.9 ( $M^+ + 1$ ).

10 Compound 1711 (Table 15 below):  $^1\text{H-NMR}$  (200MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  3.95 (d, 2H,  $J = 5.8\text{Hz}$ ), 4.23 - 4.31 (m, 2H), 4.53 (t, 2H,  $J = 8.8\text{Hz}$ ), 6.30 (d, 1H,  $J = 3.0\text{Hz}$ ), 6.98 (d, 1H,  $J = 3.0\text{Hz}$ ), 7.45 - 7.48 (m, 3H), 7.83 - 8.42 (m, 2H), 9.70 (brs, 1H). MS (ES): 281.1 ( $\text{M}^+ + 1$ ).

20

25

30

35

**Compound 1716 (Table 15 below):** <sup>1</sup>H-NMR (200MHz, CD<sub>3</sub>OD) δ 1.65 (m, 1H), 2.18 (m, 1H), 2.49 (br d, 2H, J = 6.2Hz), 2.64 (m, 1H), 3.38 (m, 1H), 3.69 (s, 3H), 3.72 (m, 1H), 3.93 (m, 1H), 4.10 (m, 1H), 5.06 (2, 2H), 6.58 (s, 1H), 6.92 (m, 2H), 7.02 (m, 1H), 7.23 (dd, 1H, J = 8.1Hz), 7.39 (m, 3H), 8.32 (m, 2H). MS (ES): 477.1 (M<sup>+</sup>+1)..

Compound 1717 (Table 15 below):  $^1\text{H}$ -NMR (200MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  1.69 (m, 1H), 2.26 (m, 1H), 2.42 (d, 2H,  $J = 7.4\text{Hz}$ ), 2.72 (m, 1H), 3.53 (m, 1H), 3.83 (m, 1H), 4.02 (m, 1H), 4.14 (dd, 1H,  $J = 10.6, 7.0\text{Hz}$ ), 5.14 (t, 2H), 6.69 (s, 1H), 6.96 (m, 2H), 7.06 (m, 1H), 7.25 (dd, 1H,  $J = 8.0\text{Hz}$ ), 7.39 (m, 3H), 8.35 (m, 2H). MS (ES): 462.2 ( $\text{M}^+ + 1$ ).

Compound 1718 (Table 15 below):  $^1\text{H}$ -NMR (200MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  1.40 - 2.00 (m, 5H), 3.52 (d, 2H, 7.6Hz), 3.80 - 4.00 (m, 1H), 4.00 - 4.20 (m, 3H), 4.50 (m, 2H), 6.36 - 6.50 (m, 2H), 6.54 (s, 1H), 6.84 - 6.92 (m, 1H), 7.05 (t, 1H,  $J = 8.2\text{Hz}$ ), 7.30 - 7.45 (m, 3H), 8.24 (d, 2H,  $J = 9.8\text{Hz}$ ). MS (ES): 449.0 ( $\text{M}^+ + 1$ ).

15

20

25

30

35

**Yeast  $\beta$ -Galactosidase reporter gene assays for human adenosine  $A_1$  and  $A_{2a}$  receptor:** Yeast strains (*S. cerevisiae*) were transformed with human adenosine  $A_1$  ( $A_1R$ ; CADUS strain CY12660) or human  $A_{2a}$  ( $A_{2a}$ ; CADUS strain CY8362) and the addition of a *lacZ*( $\beta$ -Galactosidase) reporter gene to utilize as a functional readout. A complete description of the transformations is listed below (see Yeast Strains). NECA (5'-N-ethylcarboxamidoadenosine), a potent adenosine receptor agonist with similar affinity for  $A_1$  and  $A_{2a}$  receptors, was used as a ligand for all assays. Test compounds were examined at 8 concentrations (0.1 - 10,000 nM) for ability to inhibit NECA-induced  $\beta$ -Galactosidase activity by CY12660 or CY8362.

**Preparation of Yeast Stock Cultures:** Each of the respective yeast strains, CY12660 and CY8362, were streaked onto an LT agar plate and incubated at 30°C until colonies were observed. Yeast from these colonies were added to LT liquid (pH 6.8) and grown overnight at 30°C. Each yeast strain was then diluted to an  $OD_{600} = 1.0-2.0$  (approximately  $1-2 \times 10^7$  cells/ml), as determined spectrophotometrically (Molecular Devices VMAX). For each 6 ml of yeast liquid culture, 4 ml of 40% glycerol (1:1.5 vol:vol) was added ("yeast/glycerol stock"). From this yeast/glycerol stock, ten 1 ml aliquots were prepared and stored at -80°C until required for assay.

**Yeast  $A_1$  R and  $A_{2a}$ R Assay:** One vial each of CY8362 and CY12660 yeast/glycerol stock was thawed and used to inoculate Supplemented LT liquid media, pH 6.8 (92 ml LT liquid, to which is added: 5 ml of 40% glucose, 0.45 ml of 1M KOH and 2.5 ml of Pipes, pH 6.8). Liquid cultures were grown 16-18 hr (overnight) at 30°C. Aliquots from overnight cultures were then diluted in LT media, containing 4U/ml adenosine deaminase (Type VI or VII from calf intestinal mucosa, Sigma), to obtain  $OD_{600} = 0.15$  ( $1.5 \times 10^6$  cells/ml) for CY8362 ( $A_{2a}R$ ) and  $OD_{600} = 0.50$  ( $5 \times 10^6$  cells/ml) for CY12660 ( $A_1R$ ).

Assays were conducted with a final volume of 100 ul in 96-well microtiter plates, such that a final concentration of 2% DMSO was achieved in all wells. For primary screening, 1-2 concentrations of test compounds were utilized (10 uM, 1 uM ).

5 For compound profiling, 8 concentrations were tested (10000, 1000, 500, 100, 50, 10, 1 and 0.1 nM). To each microtiter plate, 10 ul of 20% DMSO was added to "Control" and "Total" wells while 10 ul of Test Compound (in 20% DMSO) was added to "Unknown" wells. Subsequently, 10 ul of NECA (5 uM for A<sub>1</sub>R,

10 1 uM for A<sub>2a</sub>R) were added to "Total" and "Unknown" wells; 10 ul of PBS was added to the "Control" wells. In the final addition, 80 ul of yeast strain, CY8362 or CY12660, were added to all wells. All plates were then agitated briefly (LabLine orbital shaker 2-3 min) and allowed to incubate for

15 4 hrs. at 30°C in a dry oven.

β-Galactosidase activity can be quantitated using either colorimetric (e.g., ONPG, CPRG), luminescent (e.g., Galacton-Star) or fluorometric substrates (e.g., FDG, Resorufin)

20 substrates. Currently, fluorescence detection is preferred on the basis of superior signal:noise ratio, relative freedom from interference and low cost. Fluorescein digalactopyranoside (FDG, Molecular Probes or Marker Gene Technologies), a fluorescent β-Galactosidase substrate, was

25 added to all wells at 20 ul/well (final concentration = 80 uM). Plates were shaken for 5-6 sec (LabLine orbital shaker) and then incubated at 37°C for 90 min (95% O<sub>2</sub>/5% CO<sub>2</sub> incubator). At the end of the 90 min incubation period, β-Galactosidase activity was stopped using 20 ul/well of 1M

30 Na<sub>2</sub>CO<sub>3</sub> and all plates shaken for 5-6 sec. Plates were then agitated for 6 sec and relative fluorescence intensity determined using a fluorometer (Tecan Spectrafluor; excitation = 485 nm, emission = 535 nm).

35 Calculations: Relative fluorescence values for "Control" wells were interpreted as background and subtracted from "Total" and "Unknown" values. Compound profiles were analyzed via logarithmic transformation (x-axis: compound concentration)

followed by one site competition curve fitting to calculate  $IC_{50}$  values (GraphPad Prism).

**Yeast strains:** *Saccharomyces cerevisiae* strains CY12660

5 [far1\*1442 tbt1-1 fus1-HIS3 can1 stel4::trp1::LYS2 ste3\*1156  
gpa1(41)-Gxi3 lys2 ura3 leu2 trp1: his3; LEU2 PGKp-  
Mfx1Leader-hA1R-PHO5term 2mu-orig REP3 Ampr] and CY8362  
[gpalp-rGxsE10K far1\*1442 tbt1-1 fus1-HIS3 can1 stel4::trp1:  
LYS2 ste3\*1156 lys2 ura3 leu2 trp1 his3; LEU2 PGKp-hA2aR 2mu-  
10 ori REP3 Ampr] were developed.

**LT Media:** LT (Leu-Trp supplemented) media is composed of 100g  
DIFCO yeast nitrogen base, supplemented with the following:  
1.0g valine, 1.0g aspartic acid, 0.75g phenylalanine, 0.9g  
15 lysine, 0.45g tyrosine, 0.45g isoleucine, 0.3g methionine,  
0.6g adenine, 0.4g uracil, 0.3g serine, 0.3g proline, 0.3g  
cysteine, 0.3g arginine, 0.9g histidine and 1.0g threonine.

## 20 Construction of Yeast Strains Expressing Human $A_1$ Adenosine Receptor

In this example, the construction of yeast strains expressing  
a human  $A_1$  adenosine receptor functionally integrated into the  
yeast pheromone system pathway is described.

25

### I. Expression Vector Construction

To construct a yeast expression vector for the human  $A_1$   
adenosine receptor, the  $A_1$  adenosine receptor cDNA was  
obtained by reverse transcriptase PCR of human hippocampus  
30 mRNA using primers designed based on the published sequence  
of the human  $A_1$  adenosine receptor and standard techniques.  
The PCR product was subcloned into the NcoI and XbaI sites of  
the yeast expression plasmid pMP15.

35 The pMP15 plasmid was created from pLPXt as follows: The  
XbaI site of YEP51 (Broach, J.R. et al. (1983) "Vectors for



high-level, inducible expression of cloned genes in yeast" p. 83-117 in M. Inouye (ed.), Experimental Manipulation of Gene Expression. Academic Press, New York) was eliminated by digestion, end-fill and religation to create Yep51NcoDXba.

5 Another XbaI site was created at the BamHI site by digestion with BamHI, end-fill, linker (New England Biolabs, # 1081) ligation, XbaI digestion and re-ligation to generate YEP51NcoXt. This plasmid was digested with Esp31 and NcoI and ligated to Leu2 and PGKp fragments generated by PCR. The

10 2 kb Leu2 PCR product was generated by amplification from YEP51Nco using primers containing Esp31 and BglII sites. The 660 base pair PGKp PCR product was generated by amplification from pPGK $\alpha$ s (Kang, Y.-S. et al. (1990) Mol Cell Biol. 10:2582-2590) with PCR primers containing BglII and NcoI

15 sites. The resulting plasmid is called pLPXt. pLPXt was modified by inserting the coding region of the a-factor prepro leader into the NcoI site. The prepro leader was inserted so that the NcoI cloning site was maintained at the 3' end of the leader, but not regenerated at the 5' end. In

20 this way receptors can be cloned by digestion of the plasmid with NcoI and XbaI. The resulting plasmid is called pMP15.

The pMP15 plasmid into which was inserted the human A<sub>1</sub> adenosine receptor cDNA was designated p5095. In this

25 vector, the receptor cDNA is fused to the 3' end of the yeast a-factor prepro leader. During protein maturation the prepro peptide sequences are cleaved to generate mature full-length receptor. This occurs during processing of the receptor through the yeast secretory pathway. This plasmid is

30 maintained by Leu selection (i.e., growth on medium lacking leucine). The sequence of the cloned coding region was determined and found to be equivalent to that in the published literature (GenBank accession numbers S45235 and S56143).

35

## II. Yeast Strain Construction

To create a yeast strain expressing the human A<sub>1</sub> adenosine receptor, yeast strain CY7967 was used as the starting

parental strain. The genotype of CY7967 is as follows:

MAT $\alpha$  gpaD1163 gpa1(41)G $\alpha$ i3 far1D1442 tbt-1 FUS1-HIS3  
 can1 ste14::trp1::LYS2 ste3D1156 lys2 ura3 leu2 trp1  
 5 his3

The genetic markers are reviewed below:

- MAT $\alpha$ ..... Mating type  $\alpha$ .
- 10 .....  
 gpa1D1163..... The endogenous yeast G-protein GPA1 has been deleted.  
 gpa1(41)G $\alpha$ i3..... gpa1(41)-Gai3 was integrated into the yeast genome. This chimeric Ga protein is composed of the first 41 amino acids of the endogenous yeast Ga subunit GPA1 fused to the mammalian G-protein Gai3 in which the cognate N-terminal amino acids have been deleted.  
 ..  
 far1D1442..... FAR1 gene (responsible for cell cycle arrest) has been deleted (thereby preventing cell cycle arrest upon activation of the pheromone response pathway).
- 15 tbt-1..... strain with high transformation efficiency by electroporation.
- FUS1-HIS3..... a fusion between the FUS1 promoter and the HIS3 coding region (thereby creating a pheromone inducible HIS3 gene).
- can 1..... arginine/canavanine permease.  
 ste14::trp1::L gene disruption of STE14, a C-farnesyl
- 20 YS2.... methyltransferase (thereby lowering basal signaling through the pheromone pathway).  
 ste3D1156..... endogenous yeast STR, the  $\alpha$  factor  
 . pheromone receptor (STE3) was disrupted.  
 lys2..... defect in 2-aminoapitate reductase, yeast need lysine to grow.  
 ura3..... defect in orotidine-5'-phosphate decarboxylase, yeast need uracil to grow
- 25 leu2..... defect in b-isopropylmalate dehydrogenase, yeast need leucine to grow.  
 trp1..... defect in phosphoribosylanthranilate, yeast need tryptophan to grow.  
 his3..... defect in imidazoleglycerolphosphate dehydrogenase, yeast need histidine to grow.

007967 007967 007967

The resultant strain carrying p5095 and p1584, referred to as CY12660, expresses the human A<sub>1</sub> adenosine receptor. To grow  
15 this strain in liquid or on agar plates, minimal media  
lacking leucine and tryptophan was used. To perform a growth  
assay on plates (assaying FUS1-HIS3), the plates were at pH  
6.8 and contained 0.5-2.5 mM 3-amino-1,2,4-triazole and  
lacked leucine, tryptophan and histidine. As a control for  
20 specificity, a comparison with one or more other yeast-based  
seven transmembrane receptor screens was included in all  
experiments.

## Construction of Yeast Strains Expressing Human A2a Adenosine Receptor

In this example, the construction of yeast strains expressing a human A2a adenosine receptor functionally integrated into the yeast pheromone system pathway is described.

### I. Expression Vector Construction

To construct a yeast expression vector for the human A2a adenosine receptor, the human A2a receptor cDNA was obtained from Dr. Phil Murphy (NIH). Upon receipt of this clone, the A2a receptor insert was sequenced and found to be identical to the published sequence (GenBank accession # S46950). The receptor cDNA was excised from the plasmid by PCR with VENT polymerase and cloned into the plasmid pLPBX, which drives receptor expression by a constitutive Phosphoglycerate Kinase (PGK) promoter in yeast. The sequence of the entire insert was once again sequenced and found to be identical with the published sequence. However, by virtue of the cloning strategy employed there were three amino acids appended to the carboxy-terminus of the receptor, GlySerVal.

### II. Yeast Strain Construction

To create a yeast strain expressing the human A2a adenosine receptor, yeast strain CY8342 was used as the starting parental strain. The genotype of CY8342 is as follows:  
MATa far1D1442 tbt1-1 lys2 ura3 leu2 trp1 his3 fus1-HIS3 can1 ste3D1156 gpaD1163 stel4::trp1::LYS2 gpalp-rG<sub>αs</sub>E10K (or gpalp-rG<sub>αs</sub>D229S or gpalp-rG<sub>αs</sub>E10K+D229S)

The genetic markers are as described in Example 1, except for the G-protein variation. For human A2a receptor-expression, yeast strains were utilized in which the endogenous yeast G protein GPA1 had been deleted and replaced by a mammalian G<sub>αs</sub>. Three rat G<sub>αs</sub> mutants were utilized. These variants contain one or two point mutations which convert them into proteins which couple efficiently to yeast βγ. They are identified as G<sub>αs</sub>E10K (in which the glutamic acid at position ten is

replaced with lysine), G<sub>os</sub>D229S (in which the aspartic acid at position 229 is replaced with serine) and G<sub>os</sub>E10K+D229S (which contains both point mutations).

- 5 Strain CY8342 (carrying one of the three mutant rat G<sub>os</sub> proteins) was transformed with either the parental vector pLPBX (Receptor<sup>-</sup>) or with pLPBX-A2a (Receptor<sup>+</sup>). A plasmid with the FUS1 promoter fused to  $\beta$ -galactosidase coding sequences (described in above) was added to assess the  
10 magnitude of activation of the pheromone response pathway.

#### Functional Assay using Yeast Strains Expressing Human A<sub>1</sub> Adenosine Receptor

- In this example, the development of a functional screening  
15 assay in yeast for modulators of the human A<sub>1</sub> adenosine receptor is described.

#### I. Ligands Used in Assay

- Adenosine, a natural agonist for this receptor, as well as  
20 two other synthetic agonists were utilized for development of this assay. Adenosine, reported to have an EC<sub>50</sub> of approximately 75 nM, and (-)-N6-(2-phenylisopropyl)-adenosine (PIA) with a reported affinity of approximately 50 nM were used in a subset of experiments. 5'-N-ethylcarboxamido-  
25 adenosine (NECA) was used in all growth assays. To prevent signaling due to the presence of adenosine in the growth media, adenosine deaminase (4U/ml) was added to all assays.

#### II. Biological Response in Yeast

- 30 The ability of the A<sub>1</sub> adenosine receptor to functionally couple in a heterologous yeast system was assessed by introducing the A<sub>1</sub> receptor expression vector (p5095, described above) into a series of yeast strains that expressed different G protein subunits. The majority of  
35 these transformants expressed G<sub>q</sub> subunits of the G<sub>q1</sub> or G<sub>qo</sub> subtype. Additional G<sub>q</sub> proteins were also tested for the possible identification of promiscuous receptor-G<sub>q</sub> protein coupling. In various strains, a STE18 or a chimeric STE18-

Gy2 construct was integrated into the genome of the yeast. The yeast strains harbored a defective HIS3 gene and an integrated copy of FUS1-HIS3, thereby allowing for selection in selective media containing 3-amino-1,2,4-triazole (tested at 0.2, 0.5 and 1.0 mM) and lacking histidine. Transformants were isolated and monolayers were prepared on media containing 3-amino-1,2,4-triazole, 4 U/ml adenosine deaminase and lacking histidine. Five microliters of various concentrations of ligand (e.g., NECA at 0, 0.1, 1.0 and 10 mM) was applied. Growth was monitored for 2 days. Ligand-dependent growth responses were tested in this manner in the various yeast strains. The results are summarized in Table 1 below. The symbol (-) indicates that ligand-dependent receptor activation was not detected while (+) denotes ligand-dependent response. The term "LIRMA" indicates ligand independent receptor mediated activation.

Table 3

Yeast strain	G $\alpha$ subunit	G $\gamma$ subunit	Strain Variants	Result
CY1316	GPA <sub>1</sub>	STE18		-
	GPA41-G $\alpha_{11}$			+
	GPA41-G $\alpha_{12}$			+
	GPA41-G $\alpha_{13}$			+
	GPA41-G $\alpha_{11}$ -G $\alpha_{OE}$			LIRMA
	GPA41-G $\alpha_{SE10K}$			-
	GPA41-G $\alpha_{SD229S}$			-
CY7967	GPA41-G $\alpha_{13}$ -integrated	STE18		+++
	GPA <sub>1</sub>	STE18	sst2 $\Delta$	+
	GPA41-G $\alpha_{11}$			+
	GPA41-G $\alpha_{12}$			+
	GPA41-G $\alpha_{13}$			+
	GPA41-G $\alpha_{11}$ -G $\alpha_{OE}$			LIRMA
	GPA41-G $\alpha_{SE10K}$			-
CY9438	GPA <sub>1</sub>	STE18-G $\gamma$ 2		-
	GPA41-G $\alpha_{11}$			+
	GPA41-G $\alpha_{12}$			+
	GPA41-G $\alpha_{13}$			+
	GPA41-G $\alpha_{11}$ -G $\alpha_{OE}$			LIRMA
	GPA41-G $\alpha_{SE10K}$			-
	GPA41-G $\alpha_{SD229S}$			-
CY10560	GPA <sub>1</sub> -integrated	STE18-G $\gamma$ 2	sst2 $\Delta$	++

As indicated in Table 3, the most robust signaling was found to occur in a yeast strain expressing the GPA<sub>1</sub>(41)-G $\alpha_{13}$  chimera.

### III. fus1-LacZ Assay

To characterize activation of the pheromone response pathway more fully, synthesis of  $\beta$ -galactosidase through fus1LacZ in response to agonist stimulation was measured. To perform the  $\beta$ -galactosidase assay, increasing concentrations of ligand were added to mid-log culture of human A<sub>1</sub> adenosine receptor

expressed in a yeast strain co-expressing a Stel8-Gy2 chimera and GPA<sub>41</sub>-G<sub>α13</sub>. Transformants were isolated and grown overnight in the presence of histidine and 4 U/ml adenosine deaminase. After five hours of incubation with 4 U/ml adenosine deaminase and ligand, induction of β-galactosidase was measured using CPRG as the substrate for β-galactoside. 5 x 10<sup>5</sup> cells were used per assay.

The results obtained with NECA stimulation indicated that at a NECA concentration of 10<sup>-8</sup> M approximately 2-fold stimulation of β-galactosidase activity was achieved. Moreover, a stimulation index of approximately 10-fold was observed at a NECA concentration of 10<sup>-5</sup> M.

The utility of this assay was extended by validation of the activity of antagonists on this strain. Two known adenosine antagonist, XAC and DPCPX, were tested for their ability to compete against NECA (at 5 mM) for activity in the β-galactosidase assay. In these assays, β-galactosidase induction was measured using FDG as the substrate and 1.6 x 10<sup>5</sup> cells per assay. The results indicated that both XAC and DPCPX served as potent antagonists of yeast-expressed A<sub>1</sub> adenosine receptor, with IC<sub>50</sub> values of 44 nM and 49 nM, respectively.

In order to determine if this inhibitory effect was specific to the A<sub>1</sub> subtype, a series of complementary experiments were performed with the yeast-based A<sub>2a</sub> receptor assay (described in Example 4). Results obtained with the A<sub>2a</sub> yeast-based assay indicated that XAC was a relatively effective A<sub>2a</sub> receptor antagonist, consistent with published reports. In contrast, DPCPX was relatively inert at this receptor, as expected from published reports.

#### IV. Radioligand Binding

The A<sub>1</sub> adenosine receptor assay was further characterized by measurement of the receptor's radioligand binding parameters. Displacement binding of [<sup>3</sup>H]CPX by several adenosine receptor



reference compounds, XAC, DPCPX, and CGS, was analyzed using membranes prepared from yeast expressing the human A<sub>1</sub> adenosine receptor. The results with yeast membranes expressing the human A<sub>1</sub> adenosine receptor were compared to those from yeast membranes expressing the human A<sub>2a</sub> adenosine receptor or the human A<sub>3</sub> receptor to examine the specificity of binding. To perform the assay, fifty mg of membranes were incubated with 0.4 nM [<sup>3</sup>H]CPX and increasing concentrations of adenosine receptor ligands. Incubation was in 50 mM Tris-HCl, pH 7.4, 1 mM EDTA, 10 mM MgCl<sub>2</sub>, 0.25 % BSA and 2 U/ml adenosine deaminase in the presence of protease inhibitors for 60 minutes at room temperature. Binding was terminated by addition of ice-cold 50 mM Tris-HCl, pH 7.4 plus 10 mM MgCl<sub>2</sub>, followed by rapid filtration over GF/B filters previously soaked with 0.5 % polyethylenimine, using a Packard 96-well harvester. Data were analyzed by nonlinear least square curve fitting procedure using Prism 2.01 software. The IC<sub>50</sub> values obtained in this experiment are summarized in Table 4, below:

Table 4

Compound	IC <sub>50</sub> [nM]		
	hA <sub>1</sub> R	hA <sub>2a</sub> R	hA <sub>3</sub> R
XAC	6.6	11.7	53.1
DPCPX	8.5	326.4	1307.0
CGS-15943	13.1	15.8	55.5
NECA	215.5	294.9	34.9
R-PIA	67.6	678.1	23.6
IB-MECA	727.7	859.4	3.1
Alloxazine	1072.0	1934.0	8216.0

These data indicate that the reference compounds have affinities consistent with those reported in the literature. The data further indicate that the yeast-based assays are of sufficient sensitivity to discriminate receptor subtype specificity.

## Functional Assay using Yeast Strains Expressing Human A2a Adenosine Receptor

In this example, the development of a functional screening assay in yeast for modulators of the human A<sub>2</sub> adenosine receptor is described.

### I. Ligands Used in Assay

The natural ligand adenosine, as well as other thoroughly characterized and commercially available ligands were used for study of the human A2a receptor functionally expressed in yeast. Three ligands have been used in the establishment of this assay. They include:

	<u>Ligand</u>	<u>Reported K<sub>i</sub></u>	<u>Function</u>
15	Adenosine	500 nM	agonist
	5'-N-ethylcarboxamidoadenosine (NECA)	10-15 nM	agonist
	(-)-N6-(2-phenylisopropyl)-adenosine (PIA)	100-125 nM	agonist

To prevent signaling due to the presence of adenosine in the growth media, adenosine deaminase (4U/ml) was added to all assays.

### II. Biological Response in Yeast

A2a receptor agonists were tested for the capacity to stimulate the pheromone response pathway in yeast transformed with the A2a receptor expression plasmid and expressing either G<sub>αs</sub>E10K, G<sub>αs</sub>D229S or G<sub>αs</sub>E10K+D229S. The ability of ligand to stimulate the pheromone response pathway in a receptor dependent manner was indicated by an alteration in the yeast phenotype. Receptor activation modified the phenotype from histidine auxotrophy to histidine prototrophy (activation of fus1-HIS3). Three independent transformants were isolated and grown overnight in the presence of histidine. Cells were washed to remove histidine and diluted to 2 x 10<sup>6</sup> cells/ml. 5 μl of each transformant was spotted onto nonselective media (including histidine) or selective

media (1 mM AT) in the absence or presence of 4 U/ml adenosine deaminase. Plates were grown at 30 °C for 24 hours. In the presence of histidine both Receptor<sup>+</sup> (R<sup>+</sup>) and Receptor<sup>-</sup> (R<sup>-</sup>) strains were capable of growth. However, in the absence of histidine only R<sup>+</sup> cells grew. Since no ligand had been added to these plates two explanations were possible for this result. One possible interpretation was that the receptor bearing yeast were at a growth advantage due to Ligand Independent Receptor Mediated Activation (LIRMA). Alternatively the yeast could have been synthesizing the ligand adenosine. To distinguish between these two possibilities, an enzyme which degrades the ligand, adenosine deaminase (ADA), was added to the growing yeast and plates. In the presence of adenosine deaminase R<sup>+</sup> cells no longer grew in the absence of histidine, indicating that the yeast were indeed synthesizing ligand.

This interpretation was confirmed by an A2a growth assay in liquid. In this experiment R<sup>+</sup> yeast (a G<sub>αs</sub>E10K strain expressing the A2a receptor) were inoculated at three densities (1 x 10<sup>6</sup> cell/ml; 3 x 10<sup>5</sup> cells/ml; or 1 x 10<sup>5</sup> cells/ml) in the presence or absence of adenosine deaminase (4 U/ml). The stringency of the assay was enhanced with increasing concentrations (0, 0.1, 0.2 or 0.4 mM) of 3-amino-1,2,4-triazole (AT), a competitive antagonist of imidazoleglycerol-P dehydratase, the protein product of the HIS3 gene. In the presence of adenosine deaminase and 3-amino-1,2,4-triazole yeast grew less vigorously. However in the absence of 3-amino-1,2,4-triazole, adenosine deaminase had little effect. Thus adenosine deaminase itself had no direct effect upon the pheromone response pathway.

An alternative approach to measuring growth and one that can be miniaturized for high throughput screening is an A2a receptor ligand spot assay. A G<sub>αs</sub>E10K strain expressing the A2a receptor (A2aR<sup>+</sup>) or lacking the receptor (R<sup>-</sup>) was grown overnight in the presence of histidine and 4 U/ml adenosine deaminase. Cells were washed to remove histidine and diluted

to  $5 \times 10^6$  cells/ml.  $1 \times 10^6$  cells were spread onto selective plates containing 4 U/ml adenosine deaminase and 0.5 or 1.0 mM 3-amino-1,2,4-triazole (AT) and allowed to dry for 1 hour. 5  $\mu$ l of the following reagents were applied to the monolayer: 10 mM adenosine, 38.7 mM histidine, dimethylsulfoxide (DMSO), 10 mM PIA or 10 mM NECA. Cells were grown 24 hours at 30°C. The results showed that cells without receptor could only grow when histidine was added to the media. In contrast, R<sup>+</sup> cells only grew in areas where the A2a receptor ligands PIA and NECA had been spotted. Since the plates contained adenosine deaminase, the lack of growth where adenosine had been spotted confirmed that adenosine deaminase was active.

### 15 III. *fus1* LacZ Assay

To quantitate activation of the yeast mating pathway, synthesis of  $\beta$ -galactosidase through *fus1*LacZ was measured. Yeast strains expressing G<sub>as</sub>E10K, G<sub>as</sub>D229S or G<sub>as</sub>E10K+D229S were transformed with a plasmid encoding the human A2a receptor (R+) or with a plasmid lacking the receptor (R-). Transformants were isolated and grown overnight in the presence of histidine and 4 U/ml adenosine deaminase.  $1 \times 10^7$  cells were diluted to  $1 \times 10^6$  cells/ml and exposed to increasing concentrations of NECA for 4 hours, followed by determination of the  $\beta$ -galactosidase activity in the cells. The results demonstrated that essentially no  $\beta$ -galactosidase activity was detected in R- strains, whereas increasing amounts of  $\beta$ -galactosidase activity were detected in R+ strains expressing either G<sub>as</sub>E10K, G<sub>as</sub>D229S or G<sub>as</sub>E10K+D229S as the concentration of NECA increased, indicating a dose dependent increase in units of  $\beta$ -galactosidase detected in response to exposure to increased ligand concentration. This dose dependency was only observed in cells expressing the A2a receptor. Furthermore the most potent G<sub>as</sub> construct for the A2a receptor was G<sub>as</sub>E10K. The G<sub>as</sub>D229S construct was the second-most potent G<sub>as</sub> construct for the A2a receptor, while the G<sub>as</sub>E10K+D229S construct was the least potent of the three

G<sub>as</sub> constructs tested, although even the G<sub>as</sub>E10K+D229S construct stimulated readily detectable amounts of  $\beta$ -galactosidase activity.

- 5 For a further description of the assays identified, see U.S. Application Serial No. 09/088985, entitled "Functional Expression of Adenosine Receptors in Yeast", filed June 2, 1998 (Attorney Docket No. CPI-093), the entire contents of which are hereby incorporated herein by reference.

10

### Pharmacological Characterization of the Human Adenosine Receptor Subtypes

#### Material and Methods

- 15 **Materials.** [<sup>3</sup>H]-DPCPX [Cyclopentyl-1,3-dipropylxantine, 8-[dipropyl-2,3-<sup>3</sup>H(N)] (120.0 Ci/mmol); [<sup>3</sup>H]-CGS 21680, [carboxyethyl-<sup>3</sup>H (N)] (30 Ci/mmol) and [<sup>125</sup>I] -AB-MECA ([<sup>125</sup>I]-4-Aminobenzyl-5'-N-Methylcarboxamidoadenosine) (2,200 Ci/mmol) were purchased from New England Nuclear (Boston, MA). XAC (Xantine amine congener); NECA (5'-N-Ethylcarboxamidoadenosine); and IB-MECA from Research Biochemicals International (RBI, Natick, MA). The Adenosine Deaminase and Complete protease inhibitor cocktail tablets were purchased from Boehringer Mannheim Corp. (Indianapolis, IN). Membranes from HEK-293 cells stably expressing the human Adenosine 2a [RB-HA2a]; Adenosine 2b [RB-HA2b] or Adenosine 3 [RB-HA3] receptor subtypes, respectively were purchased from Receptor Biology (Beltsville, MD). Cell culture reagents were from Life Technologies (Grand Island, NY) except for serum that was from Hyclone (Logan, UT).
- 20
- 25
- 30

Yeast strains: *Saccharomyces cerevisiae* strains CY12660 [far1\*1442 tbt1-1 fus1-HIS3 can1 stel4::trp1::LYS2 ste3\*1156 gpa1(41)-Gxi3 lys2 ura3 leu2 trp1: his3; LEU2 PGKp-Mfo1Leader-hA1R-PHO5term 2mu-orig REP3 Ampr] and CY8362 [gpa1p-rGosE10K far1\*1442 tbt1-1 fus1-HIS3 can1 stel4::trp1: LYS2 ste3\*1156 lys2 ura3 leu2 trp1 his3; LEU2 PGKp-hA2aR 2mu-

35

Yeast culture: Transformed yeast were grown in Leu-Trp [LT] media (pH 5.4) supplemented with 2% glucose. For the preparation of membranes 250 ml of LT medium were inoculated with start titer of  $1-2 \times 10^6$  cells/ml from a 30 ml overnight culture and incubated at 30°C under permanent oxygenation by rotation. After 16 h growth the cells were harvested by centrifugation and membranes were prepared as described below.

Yeast Cell Membrane Preparations: 250 ml cultures were harvested after overnight incubation by centrifugation at 2,000 x g in a Sorvall RT6000 centrifuge. Cells were washed in ice-cold water, centrifuged at 4°C and the pellet was resuspended in 10 ml ice-cold lysis buffer [5 mM Tris-HCl, pH 7.5; 5 mM EDTA; and 5 mM EGTA] supplemented with Protease inhibitor cocktail tablets (1 tablet per 25 ml buffer). Glass beads (17 g; Mesh 400-600; Sigma) were added to the suspension and the cells were broken by vigorous vortexing at 4°C for 5 min. The homogenate was diluted with additional 30 ml lysis buffer plus protease inhibitors and centrifuged at 3,000 x g for 5 min. Subsequently the membranes were pelleted at 36,000 x g (Sorvall RC5B, type SS34 rotor) for 45 min. The resulting membrane pellet was resuspended in 5 ml membrane buffer [50 mM Tris-HCl, pH 7.5; 0.6 mM EDTA; and 5 mM MgCl<sub>2</sub>] supplemented with Protease inhibitor cocktail tablets (1 tablet per 50 ml buffer) and stored at -80°C for further experiments.

**Mammalian Cell Membrane Preparations:** HEK-293 cell membranes were prepared as described previously (Duzic E et al.: *J. Biol. Chem.*, 267, 9844-9851, 1992) Briefly, cells were  
5 washed with PBS and harvested with a rubber policeman. Cells were pelted at 4°C 200 x g in a Sorvall RT6000 centrifuge. The pellet was resuspended in 5 ml/dish of lysis buffer at 4°C (5 mM Tris-HCl, pH 7.5; 5 mM EDTA; 5 mM EGTA; 0.1 mM Phenylmethylsulfonyl fluoride, 10 mg/ml pepstatin A; and 10  
10 mg/ml aprotinin) and homogenized in a Dounce homogenizer. The cell lysate was then centrifuged at 36,000 x g (Sorvall RC5B, type SS34 rotor) for 45 min and the pellet resuspended in 5 ml membrane buffer [50 mM Tris-HCl, pH 7.5; 0.6 mM EDTA; 5 mM MgCl<sub>2</sub>; 0.1 mM Phenylmethylsulfonyl fluoride, 10 mg/ml  
15 pepstatin A; and 10 mg/ml aprotinin) and stored at -80 °C for further experiments.

The Bio-Rad protein assay kits, based on the Bradford dye-binding procedure, (Bradford, M.: *Anal. Biochem.* 72:248  
20 (1976)) were used to determine total protein concentration in yeast and mammalian membranes.

**Adenosine 1 receptor subtype saturation and competition radioligand binding:** Saturation and competition binding on  
25 membranes from yeast cell transformed with human A<sub>1</sub> receptor subtype were carried out using antagonist [<sup>3</sup>H] DPCPX as a radioactive ligand. Membranes was diluted in binding buffer [50 mM Tris-HCl, pH 7.4; containing 10 mM MgCl<sub>2</sub>; 1.0 mM EDTA; 0.25% BSA; 2 U/ml adenosine deaminase and 1 protease  
30 inhibitor cocktail tablet/50 ml] at concentrations of 1.0 mg/ml.

In saturation binding membranes (50 µg/well) were incubate with increasing concentrations of [<sup>3</sup>H] DPCPX (0.05 - 25 nM)  
35 in a final volume of 100 µl of binding buffer at 25°C for 1 hr in the absence and presence of 10 µM unlabeled XAC in a 96-well microtiter plate.

In competition binding membranes (50  $\mu$ g/well) were incubate with [ $^3$ H] DPCPX (1.0 nM) in a final volume of 100  $\mu$ l of binding buffer at 25°C for 1 hr in the absence and presence of 10  $\mu$ M unlabeled XAC or increasing concentrations of competing compounds in a 96-well microtiter plate.

**Adenosine 2a receptor subtype competition radioligand binding:** Competition binding on membranes from HEK293 cell stably expressing the human A2a receptor subtype were carried out using agonist [ $^3$ H] CGS-21680 as a radioactive ligand. Membranes was diluted in binding buffer [50 mM Tris-HCl, pH 7.4; containing 10 mM MgCl<sub>2</sub>; 1.0 mM EDTA; 0.25% BSA; 2 U/ml adenosine deaminase and 1 protease inhibitor cocktail tablet/50 ml] at concentrations of 0.2 mg/ml. Membranes (10  $\mu$ g/well) were incubate with [ $^3$ H] CGS-21680 (100 nM) in a final volume of 100  $\mu$ l of binding buffer at 25°C for 1 hr in the absence and presence of 50  $\mu$ M unlabeled NECA or increasing concentrations of competing compounds in a 96-well microtiter plate.

**Adenosine 3 receptor competition radioligand binding:** Competition binding on membranes from HEK293 cell stably expressing the human A3 receptor subtype were carried out using agonist [ $^{125}$ I] AB-MECA as a radioactive ligand. Membranes was diluted in binding buffer [50 mM Tris-HCl, pH 7.4; containing 10 mM MgCl<sub>2</sub>; 1.0 mM EDTA; 0.25% BSA; 2 U/ml adenosine deaminase and 1 protease inhibitor cocktail tablet/50 ml] at concentrations of 0.2 mg/ml. Membranes (10  $\mu$ g/well) were incubate with [ $^{125}$ I] AB-MECA (0.75 nM) in a final volume of 100  $\mu$ l of binding buffer at 25°C for 1 hr in the absence and presence of 10  $\mu$ M unlabeled IB-MECA or increasing concentrations of competing compounds in a 96-well microtiter plate.

At the end of the incubation, the A<sub>1</sub>, A<sub>2a</sub> and A<sub>3</sub> receptor subtypes radioligand binding assays was terminated by the addition of ice-cold 50 mM Tris-HCl (pH 7.4) buffer



supplemented with 10 mM  $MgCl_2$ , followed by rapid filtration over glass fiber filters (96-well GF/B UniFilters, Packard) previously presoaked in 0.5% polyethylenimine in a Filtermate 196 cell harvester (Packard). The filter plates were dried  
5 coated with 50  $\mu$ l /well scintillation fluid (MicroScint-20, Packard) and counted in a TopCount (Packard). Assays were performed in triplicate. Non-specific binding was  $5.6 \pm 0.5\%$ ,  $10.8 \pm 1.4\%$  and  $15.1 \pm 2.6\%$  of the total binding in a A1R, A2aR and A3R binding assay, respectively.

10

**Adenosine 2b receptor subtype competition radioligand binding:** Competition binding on membranes from HEK293 cell stably expressing the human A2b receptor subtype were carried out using A<sub>1</sub> receptor antagonist [<sup>3</sup>H] DPCPX as a radioactive  
15 ligand. Membranes was diluted in binding buffer [10 mM Hepes-KOH, pH 7.4; containing 1.0 mM EDTA; 0.1 mM Benzamidine and 2 U/ml adenosine deaminase] at concentrations of 0.3 mg/ml. Membranes (15  $\mu$ g/well) were incubate with [<sup>3</sup>H] DPCPX (15 nM) in a final volume of 100  $\mu$ l of binding buffer at 25°C for 1  
20 hr in the absence and presence of 10  $\mu$ M unlabeled XAC or increasing concentrations of competing compounds in a 96-well microtiter plate. At the end of the incubation, the assay was terminated by the addition of ice-cold 10 mM Hepes-KOH (pH 7.4) buffer followed by rapid filtration over glass fiber  
25 filters (96-well GF/C UniFilters, Packard) previously presoaked in 0.5% polyethylenimine in a Filtermate 196 cell harvester (Packard). The filter plates were dried coated with 50  $\mu$ l/well scintillation fluid (MicroScint-20, Packard) and counted in a TopCount (Packard). Assays were performed  
30 in triplicate. Non-specific binding was  $14.3 \pm 2.3\%$  of the total binding.

Specific binding of [<sup>3</sup>H] DPCPX; [<sup>3</sup>H] CGS-21680 and [<sup>125</sup>I] AB-MECA was defined as the difference between the total binding  
35 and non-specific binding. Percent inhibition of the compounds was calculated against total binding. Competition data were analyzed by iterative curve fitting to a one site model, and

007027 97552.60

$K_I$  values were calculated from  $IC_{50}$  values (Cheng and Prusof, Biochem. Pharmacol. 22, 3099-3109, 1973) using the GraphPad Prism 2.01 software.

## 5 Results

A primary function of certain cell surface receptors is to recognize appropriate ligands. Accordingly, we determined ligand binding affinities to establish the functional integrity of the Adenosine 1 receptor subtype expressed in yeast. Crude membranes prepared from *Saccharomyces cerevisiae* transformed with human Adenosine 1 receptor subtype construct exhibited specific saturable binding of [ $^3H$ ] DPCPX with a  $K_D$  of  $4.0 \pm 0.19$  nM. The  $K_D$  and  $B_{max}$  value were calculated from the saturation isotherm and Scatchard transformation of the data indicated a single class of binding sites. The densities of adenosine binding sites in the yeast membrane preparations were estimated to  $716.8 \pm 43.4$  fmol/mg membrane protein.

The pharmacological subtype characteristics of the recombinant yeast cells transformed with human  $A_1$  receptor subtype were investigated with subtype selective adenosine ligands (XAC, DPCPX; CGS-15943; Compound 600; Compound 1002; NECA, (R)-PIA; IB-MECA and Alloxazine) that competed with [ $^3H$ ] DPCPX in the expected rank order. Displacement curves recorded with these compounds show the typical steepness with all the ligands, and the data for each of the ligands could be modeled by a one-site fit. The apparent dissociation constants estimated for the individual compound from the curves (Table 5) are consistent with value published for the receptor obtained from other sources.

Table 5


K<sub>i</sub> values for membranes from yeast cells transformed with  
5 human A<sub>1</sub> receptor subtype

Ligands	K <sub>i</sub> (nM)
XAC	5.5
DPCPX	7.1
CGS-1594	10.8
NECA	179.6
15 (R)-PIA	56.3
IB-MECA	606.5
Alloxazine	894.1
Compound 600	13.9
20 Compound 1002	9.8









Tables 6 through 12 demonstrate the efficacy and structure  
activity profiles of deazapurines of the invention. Tables  
13 and 14 demonstrate selectivity can be achieved for human  
25 adenosine receptor sites by modulation of the functionality  
about the deazapurine structure. Table 14 also demonstrates  
the surprising discovery that the compounds set forth therein  
have subnanomolar activity and higher selectivity for the A<sub>2b</sub>  
receptor as compared to the compounds in Table 13.

30

### Effect of N<sub>6</sub>-Substituent



		A1	
Compound	R	Binding Ki (nM)	Yeast IC50 (nM)
600		13.9	97.2
601		1423	>10.000
602		483.5	>10.000
603		196.6	4442.0
604		>10.000	>10000
605		>10000	>10000
606		297.9	>10000

607		309.7	>10000
608		29.1	
609		193.9	
610		411.5	
611		785.6	>10000
612		64.8	
613		6726.0	
614		32.1	




615	 (dl)	816.9	2577.0
616		34.3	

TABLE 7

Effect of C<sub>2</sub>-Substituent






Compound	R	A1	
		Binding K <sub>i</sub> (nM)	Yeast IC <sub>50</sub> (nM)
700		604.5	>10000
701		157.7	763.1



TABLE 8

Effect of Pyrrole Ring Substituent



Compound	R	R'	R''	R'''	A1	
					Binding Ki (nM)	Yeast IC50 (nM)
800		Me	Me	Me	3311	>10000
801		H	Me	H	22.3	148.3
802		H	H	Me	8.9	
803			Me	Me	2210	>10000
804			Me	Me	863.1	
805			Me	Me	4512	








806			Me	Me	8451	
807			Me	Me	35.3	

TABLE 9








		A1	
Compound	R	Binding Ki (nM)	Yeast IC50 (nM)
900		863.1	
901		4512	
902		8451	
903		35.3	

TABLE 10

Effect of N<sub>6</sub>-Substituent



Compound	R	A1	
		Binding K <sub>i</sub> (nM)	Yeast IC <sub>50</sub> (nM)
1000		1789	>10000
1001		54.4	1865
1002		9.8	82.8
1003		26.7	195.7
1004		32.8	545.8














1005		147.5	3972
1006		151.7	2918
1007		692.5	>10000
1008		93.1	3217
1009		475.3	>10000
1010		674.9	9376.0
1011		121.9	2067.5
1012		233.9	3462
1013		270.1	3009.5
1014		384.9	2005
1015		179.3	3712
1016		176.1	5054

TABLE 11

Effect of N<sub>6</sub>-Substituent



Compound	R	A1	
		Binding K <sub>i</sub> (nM)	Yeast IC <sub>50</sub> (nM)
1100		9.8	115.4
1101		53.9	551.0
1102		10.3	101.3
1103		71.1	3217
1104		6.5	58.7
1105		105.4	472.1









1106		27.8	162.4
1107		126.5	1297.0
1108		2.3	
1109		9.0	
1110		17.3	
1111		2.5	
1112		213	

TABLE 12  
"Retro-Amide" Analogues



Compound	R	A1	
		Binding Ki (nM)	Yeast IC50 (nM)
1200		16.5	189.4
1201		7.4	45.7
1202		95.8	3345.0
1203		529.1	4040.0
1204		1060.0	>10000

000000-000000








1205		1272	>10000
1206		50.8	4028
1207		48.5	701.5

TABLE 13  
Profile of Selective Adenosine Antagonists

		Binding Ki (nM)			
Compound	R	A1	A2a	A2b	A3
1300		9.8- 25.1	18.0- 48.6	80.3	513.0
1301		27.8	50.7	84.6	429.8
1302		20.2	75.6	20.1	4.3

001027-9796260












1309		40	109	109	0.3
1310		255	76%	275	≤2.6
1311		531	981	736	5.3
1312		443	2965	375	≤6.2
1313 <sup>3</sup>		30%	65%	515	24
1314		87	204	30	0.02

Table 14:  
Profile of Selective A<sub>2c</sub> Antagonists

5

10



15




20

25

Compound	XR <sub>1</sub>	R <sub>2</sub>	Binding Data K <sub>i</sub> (nM)			
			A <sub>1</sub>	A <sub>2a</sub>	A <sub>2B</sub>	A <sub>3</sub>
1400	-O-Ph	Me	41.7	21	10.3	14.6
1401	-O-Ph(p)F	Me	33	58	8.8	18
1402	-O-Ph(p)Cl	Me	825	591	22	60
1403	-N-pyridin-2-one	Me	60	41	18	48
1404	-NH-Ph	Me	49	31	4.6	57




TABLE 15. Adenosine A<sub>3</sub> Receptor Selective Compounds

\* at least 10 times more selective than other three subtypes.

Compound	Structure	Ki-A1	Ki-A2a	Ki-A2b	Ki-A3
1202					*
1700					*
1309					*








1703					*
1704					*
1705					*





2025-06-16 16:33:40



1709					*
1710					*
1711					*


000000 97932600



1712					*
1713					*
1714					*
1715					*


1716					*
1717					*
1718					*

1719



✱

This invention also provides a compound having the structure:



1720

In a further embodiment the invention provides a method for inhibiting the activity of an A<sub>3</sub> adenosine receptor in a cell, which comprises contacting the cell with the compound 1720.

In a further embodiment the invention provides a method for inhibiting the activity of an A<sub>3</sub> adenosine receptor in a cell, wherein the compound is an antagonist of the A<sub>3</sub> adenosine receptor.

In a further embodiment the invention provides the above method for inhibiting the activity of an A<sub>3</sub> adenosine receptor in a cell, wherein the cell is human cell.

In a further embodiment the invention provides the above method for inhibiting the activity of an A<sub>3</sub> adenosine receptor in a cell, wherein the cell is a human cell and wherein the compound is an antagonist of A<sub>3</sub> adenosine receptors.

In a further embodiment the invention provides a method of treating damage to the eye of a subject which comprises administering to the subject a composition comprising a therapeutically effective amount of the compound 1720.

In a further embodiment the invention provides the above method, wherein the damage comprises retinal or optic nerve head damage.

5 In a further embodiment the invention provides a therapy for glaucoma, comprising administering to a subject a therapeutically effective amount of the compound 1720.

10 In a further embodiment the invention provides a therapy for glaucoma comprising one or more adenosine receptor antagonists, preferably comprising an adenosine receptor A3 antagonist (preferably an N-6 substituted 7-deazapurine, most preferably [2-(3H-Imidazol-4-yl)-ethyl]-(2-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine).

15 In an alternative embodiment the invention provides a combination therapy for glaucoma comprising an adenosine receptor A3 antagonist (preferably an N-6 substituted 7-deazapurine, most preferably [2-(3H-Imidazol-4-yl)-ethyl]-(2-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine)) and one or more  
20 other compounds selected from the group consisting of beta adrenoceptor antagonists (i.e. beta adrenergic antagonists or b-blockers) (e.g. timolol maleate, betaxolol, carteolol, levobunolol, metipranolol, L-653328 (the acetate ester of L-652698), beta 1 adrenoceptor antagonists), alpha-2 adrenoceptor  
25 agonists (e.g. aplaclonidine, brimonidine, AGN-195795, AGN-190837 (an analog of Bay-a-6781)), carbonic anhydrase inhibitors (brinzolamide, dorzolamide, MK-927 (an inhibitor of the human carbonic anhydrase II isoenzyme), inhibitors of  
30 carbonic anhydrase IV isoenzyme), cholinergic agonists (e.g. muscarinic cholinergic agonists, carbachol, pilocarpine HCl, pilocarpine nitrate, pilocarpine, pilocarpine prodrugs (e.g. DD-22A)), prostaglandins and prostaglandin receptor agonists

(e.g. latanoprost, unoprostone isopropyl, PGF2 alpha agonists, prostanoid-selective FP receptor agonists, PG agonists such as the hypotensive prostamides), angiotensin converting enzyme (ACE) inhibitors (e.g. Spirapril, spiraprilat), AMPA receptor antagonists, 5-HT agonists (e.g. a selective 5-HT 1A receptor agonist such as MKC-242 (5-3-[(2S)-1,4-benzodioxan-2-ylmethyl)amino]propoxy-1,3-benzodioxole HCl), angiogenesis inhibitors (e.g. the steroid anecortave), NMDA antagonists (e.g. HU-211, memantine, the cannabinoid NMDA-receptor agonist dextranabinol, prodrugs and analogs of dextranabinol, NR2B-selective antagonists (e.g. eliprodil (SL-82.0715)), renin inhibitors (e.g. CGP-38560, SR-43845), cannabinoid receptor agonists (e.g. tetrahydrocannabinol (THC) and THC analogs, selective CB2 cannabinoid receptor agonists (e.g. L-768242, L-759787), compounds such as anandamide that bind to both brain-specific CB1 receptors and peripheral CB2 receptors), angiotensin receptor antagonists (e.g., angiotensin II receptor antagonists (e.g. CS-088), selective angiotensin II AT-I receptor antagonists, such as losartan potassium), hydrochlorothiazide (HCTZ), somatostatin agonists (e.g. the non-peptide somatostatin agonist NNC-26-9100), glucocorticoid antagonists, mast cell degranulation inhibitors (e.g. nedocromil), alpha-adrenergic receptor blockers (e.g. dapiprazole, alpha-2 adrenoceptor antagonists, alpha 1 adrenoceptor antagonists (e.g. bunazosin)), alpha-2 adrenoceptor antagonists, thromboxane A2 mimetics, protein kinase inhibitors (e.g. H7), prostaglandin F derivatives (e.g. S-1033), prostaglandin-2 alpha antagonists (e.g. PhXA-34), dopamine D1 and 5-HT2 agonists (fenoldopam), nitric-oxide-releasing agents (e.g. NCX-904 or NCX-905, nitric-oxide-releasing derivatives of timolol), 5-HT 2 antagonists (e.g. sarpogrelate), NMDA antagonists (e.g. prodrugs and analogs of dextranabinol), alpha 1 adrenoceptor antagonists (e.g.

bunazosin), cyclooxygenase inhibitors (e.g. diclofenac, or the non-steroidal compound nepafenac), inosine, dopamine D2 receptor and alpha 2 adrenoceptor agonists (e.g. talipexole), dopamine D1 receptor antagonist and D2 receptor agonists (e.g. SDZ-GLC-756), vasopressin receptor antagonists (e.g. vasopressin V2 receptor antagonists (e.g. SR-121463)), endothelin antagonists (e.g. TBC-2576), 1-(3-hydroxy-2-phosphonylmethoxypropyl)cytosine (HPMPC) and related analogs and prodrugs, thyroid hormone receptor ligands (e.g. KB-130015), muscarinic M1 agonists, NMDA-receptor antagonists (e.g. the cannabinoid NMDA-receptor antagonist dextranabinol), PG agonists such as the hypotensive lipids, prostamides, sodium channel blockers, NMDA antagonists, mixed-action ion channel blockers, beta adrenoceptor antagonist and PGF2 alpha agonist combinations (e.g. latanoprost and timolol), guanylate cyclase activators (e.g. atrial natriuretic peptide (ANP) or non-peptide mimetics, inhibitors of ANP neutral endopeptidase, nitrovasodilators (e.g. nitroglycerin, hydralazine, sodium nitroprusside), endothelin receptor modulators (e.g. ET-1 or non-peptide mimetics, sarafotoxin-S6c), ethacrynic acid, other phenoxyacetic acid analogs (e.g. indacrinone, ticrynafen), actin disrupters (e.g. latrunculin), calcium channel blockers (e.g. verapamil, nifedipine, brovincamine, nivaldipine) and neuroprotective agents.

A combination therapy for glaucoma, comprising the compound of 1702, and one or more compounds selected from the group consisting of beta adrenoceptor antagonists, alpha-2 adrenoceptor agonists, carbonic anhydrase inhibitors, cholinergic agonists and prostaglandin receptor agonists.

In a further embodiment the invention provides a pharmaceutical composition comprising a therapeutically effective amount of

the compound 1720 and a pharmaceutically acceptable carrier.

In a further embodiment the invention provides a packaged pharmaceutical composition for treating a disease associated with A<sub>3</sub> adenosine receptor in a subject, comprising:

(a) a container holding a therapeutically effective amount of the compound 1720; and

(b) instructions for using said compound for treating said disease in a subject.

In a further embodiment the invention provides a method of making a composition which comprises the compound 1720, the method comprising admixing the compound 1702 with a suitable carrier.


In a further embodiment the invention provides a pharmaceutically acceptable salt of compound 1720, wherein the pharmaceutically acceptable salt contains an anion selected from the group consisting of maleic, fumaric, tartaric, acetate, phosphate and mesylate.



Exemplification

**Example 37:** Synthesis of [2-(3*H*-Imidazol-4-yl)-ethyl]-(2-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)-amine (1720)

Compound **1720** was synthesized using precursor compound **1** of synthesis scheme VII to obtain:



Aryl chloride **1** (400mg, 1.50mmol), DMSO (10mL) and histamine (1.67g, 15.0mmol) are combined and heated to 120°C under nitrogen. After 6.5h, the reaction is cooled to room temperature and partitioned between EtOAc and water. The layers are separated and the aqueous layer is extracted with EtOAc (3x). The combined organic layers are washed with brine (2x), dried over MgSO<sub>4</sub>, filtered and concentrated to yield 494mg of a brown solid. The solid is washed with cold MeOH and recrystallized from MeOH to yield 197mg (43%) of an off white solid (**1720**). <sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ 3.05 (t, 2H, J = 7.0Hz), 3.94 (t, 2H, J = 7.0Hz), 6.50 (d, 1H, J = 3.5Hz), 6.88 (brs, 1H), 7.04 (d, 1H, J = 3.5Hz), 7.42 (m, 3H), 7.57 (s, 1H), 8.34 (m, 2H); MS (ES): 305.1 (M<sup>+</sup>+1); Mpt = 234-235°C.

Activity of Compounds

Adenosine 3 ( $A_3$ ) receptor competition radio ligand binding was carried out for compound 1720 as described herein and *inter alia*, on pages 167-168 of this specification. Compound 1720 was found to have an  $A_3$  receptor binding affinity greater than 10 times that of reference compound 1308 as described herein and, *inter alia*, in Table 13, on page 183 of the specification.

**Incorporation by Reference**

All patents, published patent applications and other references disclosed herein are hereby expressly incorporated herein by reference.

**Equivalents**

Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, many equivalents to specific embodiments of the invention described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.